ACADEMICIA: An International Multidisciplinary Research Journal
ISSN: 2249-7137 Vol. 12, Issue 1, January 2022 SJIF 2021 = 7.492
A peer reviewed journal

ACADEMICIA
An International Multidisciplinary Research Journal

ISSN (online) : 2249 –7137

Editor-in-Chief : Dr. B.S. Rai

Impact Factor : SJIF 2021 = 7.492

Frequency : Monthly

Country : India

Language : English

Start Year : 2011

Indexed/ Abstracted : Scientific Journal Impact Factor (SJIF2020 - 7.13), Google Scholar, CNKI Scholar, EBSCO Discovery, Summon (ProQuest), Primo and Primo Central, I2OR, ESJI, IIJIF, DRJI, Indian Science and ISRA-JIF and Global Impact Factor 2019 - 0.682

E-mail id: saarjjournal@gmail.com

VISION

The vision of the journals is to provide an academic platform to scholars all over the world to publish their novel, original, empirical and high quality research work. It propose to encourage research relating to latest trends and practices in international business, finance, banking, service marketing, human resource management, corporate governance, social responsibility and emerging paradigms in allied areas of management including social sciences , education and information & technology. It intends to reach the researcher's with plethora of knowledge to generate a pool of research content and propose problem solving models to address the current and emerging issues at the national and international level. Further, it aims to share and disseminate the empirical research findings with academia, industry, policy makers, and consultants with an approach to incorporate the research recommendations for the benefit of one and all.

WEAK FERROMAGNETISM

Fayziev Shakhobiddin Shavkatovich*; Yuldasheva Nilufar Bakhtiyorovna; Saidjonova Madinabonu Shuhratovna*****

*Candidate of Science of Physics and Mathematics,
Bukhara State University, Bukhara, UZBEKISTAN
Email id: Fayziyev_83@mail.ru

**Lecturer of Department of Physics,
Bukhara State University, Bukhara, UZBEKISTAN
Email id: nilufar.yuldasheva1995@mail.ru

***Department of Physics Masters,
Bukhara State University, Bukhara, UZBEKISTAN

DOI: 10.5958/2249-7137.2022.00054.4

ABSTRACT

The magneto optic properties of FeBO_3are determined by the limits in the dielectric absorption tensor, which depend on the ferro and antıferromagnetism vectors. A classic example of the effect of crystal magnetic symmetry on the formation of a weak ferromagnetic moment is hematite α= 〖Fe〗 _2 O_3, which has a rhombohedral elementary cell with four iron atoms. Hence the term "parasitic ferromegnetism", a term still found in some foreign literature.

KEYWORDS: *Antıferromagnetic, Hematite, Spinorientation, Crystal, Ferromagnetic Moment*

INTRODUCTION

In many antiferromagnets, the properties of the crystal structure are such that the sub-lattice atoms with opposite directions of magnetization are in less crystallographic states, so they are affected by different anisotropic forces. This (if crystal symmetry allows) can lead to non-collenear sub-lattice magnetization, their exact mutual compensation is violated, and in total the nominal $10^{-2} \div 10^{-5}$a small spontaneous magnetization occurs. Noncolenear anisotropy may be associated with both indirect exchange interactions and with single-ion anisotropy. The phenomenon of such a small spontaneous magnetization is called weak ferromagnetism, and the substances observed in this phenomenon are called weak ferromagnets. **[1-8]**

The presence of insignificant spontaneous magnetization in hematite ($\alpha - Fe_2O_3$) and a number of 3d-metal salts has long been known, and many researchers have suggested the presence of ferromagnetic (compound, foreign) primes in samples. Hence the term "parasitic ferromegnetism", a term still found in some foreign literature. During the preparation of more pure crystals, it became clear that the presence of spontaneous magnetization was a property of the crystals themselves, and that A.S. Borovik-Romanov had suggested that it was due to the noncollinearity of the bottom of the magnetic grille. The theoretical basis of this idea was given by I.E. Dzolyashinsky **[9, 10,11,12].**

In the example of rhombohedral carbonates $MnCO_3$ and $CoCO_3$belonging to the spatial group $D_{3d}^6(R3c)$and having the following elements of symmetry in the paramagnetic state, we see the formation of weak ferromagnetism: $2C_3$, $3U_2$, I, $2S_6$, $3\sigma_d$, R.

$2C_3$- third-order axes, parallel to [111];

$3U_2$- second-order axes, perpendicular to [111];

I - Inversion;

S_6 – is the axis of rotation of the sixth order mirror;

σ_d – is the translational sliding plane along the [111] axis and perpendicular to the U_2 axis.

Figure 2.2.1 shows the location of magnetic ions and some elements of symmetry in the structure of carbonates. The magnetic properties of the carbonates are well represented by the two magnetic grid substrates, and the magnetic element cell is compatible with the crystal chemistry.

Thus, the magnetic atoms 1 and 2 (Figure 2.2.1) correspond to a single elementary cell but under different magnetic grids. Some of the elements of symmetry listed are missing.

C_3: $1\rightarrow 1$, $2\rightarrow 2$, U_2: $1\leftrightarrows 2$; S_6va$I$: $1\rightarrow 1$, $2\rightarrow 2$; σ_d: $1\leftrightarrows 2$.

There are three possible types of magnetic structures (Figure 2.2.2).

In structure A, the magnetic moments under the grid are oriented along the axis [111], in structure B the magnitude σ_d lies in the plane of symmetry, and in structure C the direction σ_d is oriented along the axis. $FeCO_3$ has a structure A, while $MnCO_3$ and $CoCO_3$ have a structure B.



(Figure 2.2.1)

Figure 2.2.1.Location of magnetic (1, 2) ions along the axis in the crystals of carbonates $MnCO_3$, $FeCO_3$ and $CoCO_3$[111].

c is the point of intersection of the axis of symmetry $U_2$with the axis c- [111];

The points of intersection of the return planes for the d- [111] axis and the S_6 axis.

Structure A has all the elements of spatial crystallographic symmetry except R. Well, in such a magnetic structure [111], the resultant magnetic moment$\vec{m} = \vec{m}_\perp + \vec{m}_{||}$, which is parallel and perpendicular to the axis, is $\vec{m}_{||}$and $\vec{m}_{\perp}$, respectively. 'ladi. $\vec{m}$is invariant with respect to the C_3 axis, and $\vec{m}_A$is not invariant, so m can be invariant with respect to this symmetry operation only when$\vec{m} = 0$. Thus, there can be no weak ferromagnetism in structure A, in fact, $FeCO_3$ does not have a weak magnetic moment.



(Figure 2.2.2)

Figure 2.2.2. The 3d-metal carbonates $FeCO_3$, $MnCO_3$ and $CoCO_3$are the three possible orientations of the magnetization of the magnetic substrate.

In structure B, the magnetic moments are in the $\sigma_d$plane (perpendicular to the plane of Figure 2.2.2) and the symmetry elements are *I*, U_2, $\sigma_d$which allows the magnetic moments of the 1- and 2-ions to rotate out of the σ_d plane against each other. the moment U_2 becomes m ≠ 0 along the axis (see Figure 2.2.3). Indeed, $MnCO_3$ and $CoCO_3$compounds are weak ferromagnets. A classic example of the effect of crystal magnetic symmetry on the formation of a weak ferromagnetic moment is hematite α=Fe_2O_3, which has a rhombohedral elementary cell with four iron atoms. At a temperature of 250 K, a spin-oriented spatial transition in hematite may be appropriate, i.e., the magnetic structure changes. At a temperature T <250 K, the magnetic moments of the iron ion lie parallel to the axis of the crystal [111], and at a temperature T> 250 K, it lies in the base plane near the Neel point (950 K). In the first case, weak ferromagnetism is prohibited, while in the second case, the magnetic structure allows it. In fact, small spontaneous magnetization is observed only in the temperature range of 250K ÷ 950K. **[13-24]**

CONCLUSION

The magneto optic properties of $FeBO_3$ are determined by the limits in the dielectric absorption tensor, which depend on the vectors of ferro and antiferromagnetism, in which the magnetoooptic properties of the crystalit is sufficient to consider only the dielectric absorption tensor component, which represents the cross section of the optical index of the plane perpendicular to the C_3 axis, to analyze its properties. Studies have shown that the existing oriented spatial transitions theory $Tb_{0,2}Y_{2,8}Fe_5O_{12}$ cannot represent the multidimensional temperature-dependent evolution observed in the domain structure of the sample, and the crystallographic mechanical stress of the sample is significantly different from the spin reorientation. 'It's a mystery. **[25-28]**

REFERENCES

1. Fedorov YuM, Leksikov AA, Aksenov AE. Photo-induced optical anisotropy in iron borate. JETF. 1985;89:2099–2112.

2. Boidedaev SR, Dzhuraev DR, Sokolov BYu, Faiziev ShSh. Effect of the Transformation of the magnetic structure of a FeBO3:Mg crystal on its magnetooptical anisotropy. Optics and Spectroscopy. 2009;4(107):651-654.

3. Fayziev ShSh. Effect of light on the modulation of the magnetic order of the FeBO3:Mg crystal. Young Science 2017;(28):8-11.

4. Sharipov MZ, Sokolov BYu, Faiziev ShSh, Mirzhonova NN. Effect of rearrangement of the magnetic structure of a FeBO3: Mg crystal on its magneto-optical anisotropy. Science, technology and education. 2015;4(10):15-18.

5. Dzhuraev DR, Sokolov BYu, Faiziev ShSh. Photoinduced changes inthe space-modulated magnetic order of a FeBO3:Mg single crystal. Russian Physics Journal. 2011;3(54):382-385.

6. Dzhuraev DR, Fayziev Sh, Sokolov BYu. 'Magnetic ripple' in rhombohedral FeBO3:Mg crystal. Proceedings of fourth international conference dedicated to the eightieth anniversary of academician M.S. Saidov 'Fundamental and applied problems of physics'. 2010; pp. 342-344.

7. Valiyev UV, Saidov KS, Lukina MM. On the nature of the Faraday effect in rare earth orthoaluminate TbAlO3. Solid state physics. 1999;(11):2047-2052.

8. Dzhuraev DR, Niyazov LN, Sokolov BY. Modulated magnetic phase of structurally heterogeneous easy-plane weak ferromagnets. Technical Physics 2016;61(6):883-886.

9. Boydedaev SR, Sokolov DY, Dzhuraev DR, Fayziev Sh. The 'magnetic ripple' state in weak ferromagnetic FeBO3: Mg. Uzbekiston Fizika Zhurnali. 2009;(5):376-383

10. Valiev UV, Dzhuraev DR, Malyshev EE, Saidov KS. Electronic structure of the ground multiplet of the Dy3+ ion in the DyAlO3 orthoaluminate. OpticsandSpectroscopy. 1999;(86): 703-706

11. Valiev UV, Lukina MM, Saidov KS. On the nature of the Faraday effect in the rare-earth ortho-aluminate TbAlO3. Physics of the Solid State. 1999;(41):1880-1884

12. Djuraev DR, Niyazov LN, Saidov KS, Sokolov BYu. Investigation of the spontaneous spin-flip phase transition in terbiumyttrium iron-garnet by the magnetooptic method. Ukrainian journal of physics. 2012;(5):531-537.

13. Dzhuraev DR, Mirzhanova N, Niyazov LN, Saidov KS. The superconductivity and nanotechnology. Modern problems of semiconductor physics. 2011, pp. 125-127

14. Dzhuraev DR, Niyazov LN, Saidov KS, Sokolov BYu, Khaydarova L. The spontaneous orientation phase transition in terbium-yttrium ferrite-garnet. Modern problems of semiconductor physics. 2011. pp. 76-78

15. Djuraev DR, Niyazov LN, Saidov KS, Sokolov BYu. Changing the cubic ferrimagnetic domain structure in temperature region of spin flip transition. Uzbekiston Fizika Zhurnali. 2011;(5):359-366.

16. Valiev UV, Gruber JB, Rustamov UR, Saidov KS, Sokolov VYu. A magnetooptical study of the odd crystal field component in a terbium-yttrium aluminum garnet. Technical Physics Letters. 2003;(11):882-885.

17. Valiev UV, Dzhuraev DR, Nekvasil V, Saidov KS. Magnetooptical studies of excited states of rare-earth ions in Y3Al5O12:Tb3+. Magnetic and Superconducting Materials: Volume 2. 2000. pp. 1059-1064

18. Fayziev Sh. Investigation of the magnetic structure of FeBO3:Mg. III international scientific conference of young researchers. 2019;(1):105-107

19. Fayziev ShSh, Saidov KS, Sulaymonov ShB. Magnetic properties of rare earth garnets. Academy. 2021;4(67):4-7

20. Sharipov MZ, Faiziev ShSh. Nizomov ShK. Features of the magneto-optical properties of a single crystal of iron borate. Science, technology and education. 2021;2-2(77):5-9.

21. Bakhtiyorovna YN, Shavkatovich FSh. Modulated magnetic structures and models of their theoretical expression. Academicia: an international multidisciplinary research journal. 2021; 1(11):1172-1175

22. Fayziev ShSh, Saidov KS. Nizomova ShK. Electronic structure of the main multiplet of the dysprosium ion in orthoaluminate. Academy. 2020;11(62):4-6.

23. Fayziev ShSh, Saidov KS, Askarov MA. Dependence of the magnetically modulated structure on the field orientation in the FeBO3:Mg crystal. Bulletin of science and education. 2020;18-2(96):6-9.

24. Shavkatovich ShF, Baxtierovna NY. Changes occuring in ferromagnets by adding some mixture. Scientific reports of Bukhara State University. 2020;1(4):8-13.

25. Sharipov MZ, Sokolov BYu, Faiziev ShSh, Mirzhonova NN. Effect of rearrangement of the magnetic structure of a FeBO3: Mg crystal on its magneto-optical anisotropy. Science, technology and education. 2015;4(10):15-18.

26. Fayziev ShSh, Juraeva LR. On the magnetic properties of iron borate doped with magnesium. Modern innovations in science and technology. 2014. P. 264-266

27. Dzhuraev DR, Sokolov BYu, Faiziev ShSh. Photoinduced changes inthe space-modulated magnetic order of a FeBO3:Mg single crystal. Russian Physics Journal. 2011;3(54):382-385.

28. Boidedaev SR, Dzhuraev DR, Sokolov BYu, Faiziev ShSh. Effect of the Transformation of the magnetic structure of a FeBO3:Mg crystal on its magnetooptical anisotropy. Optics and Spectroscopy. 2009;4(107):651-654.