

The Properties of Cotton Resistance and Adaptability to Drought Stress

Kholliyev Askar Ergashovich¹, Norboyeva Umida Toshtemirovna², Adizova Khamida Raximovna³, Fayzieva Flora Abdullaevna⁴

¹Doctor of Biological Sciences, Professor, Bukhara State University, Bukhara, Uzbekistan. E-mail: a.e.holliyev@buxdu.uz

²Doctor of Biological Sciences, Professor, Bukhara State University, Bukhara, Uzbekistan. E-mail: u.t.norboyeva@buxdu.uz

³Lecturer, Department of Ecology, Bukhara State University, Bukhara, Uzbekistan. E-mail: x.r.adizova@buxdu.uz

⁴Lecturer, Department of Ecology, Bukhara State University, Bukhara, Uzbekistan. E-mail: f.a.fayzieva@buxdu.uz

Abstract

The article presents data obtained based on the study of physiological and biochemical processes of drought tolerance of cotton varieties and physiological indicators of protective adaptation. The study aims to determine the degree and form of protective adaptation of cotton varieties to soil drought based on the physiological and biochemical properties of water exchange, as well as to develop methods to increase drought tolerance. To obtain a high and high-quality yield in cotton farms located in the middle and lower regions of the Zarafshan oasis (Uzbekistan) and in areas where drought is observed, drought-resistant and adaptive features, the possibility of growing varieties of Bukhara-6, Bukhara-102, Bukhara-8 cotton with high yield and quality, as well as conclusions, suggestions and recommendations.

Keywords: Cotton, Drought, Stress, Diffusion Resistance, Transpiration, Metabolic and Bound Water, The Cell Sap, Water Deficiency, Endurance, Adaptation.

DOI: 10.47750/pnr.2022.13.04.129

INTRODUCTION

Global climate change is causing an increase in air temperature in the biosphere, and hot winds caused by a sharp drop in relative humidity in the summer months are causing atmospheric and soil drought. In the current period of serious water problems, it is important to introduce water-saving agro-technologies, as well as to develop methods of growing cotton varieties that are resistant to soil and atmospheric drought and have a high efficiency of water use [1,2].

The strongest negative impact of adverse environmental factors, such as the atmosphere and soil drought, falls on the water-demanding-critical period of cotton, i.e. the flowering stage. At the same time, due to the lack of water in the soil and high air temperatures together adversely affecting the physiological and biochemical processes that take place in the cotton, the yield and its quality decrease. Therefore, it is important to zoning cotton varieties that are resistant to such adverse factors based on specific soil and climatic conditions [3,4,5].

The negative impact of drought can be reduced to some extent by providing cotton varieties with sufficient mineral fertilizers, timely agro-technical processing, and the organization of crop rotation. It is also possible to increase the resistance of plants to the adverse effects of adverse factors by applying the electrification method [6,7].

The urgency of the above problem is that the soil and

climatic conditions of the cotton-growing areas in our country are very different. The potential of cotton varieties planted in a particular ecological zone also varies depending on the level of agro-technical processing. One of the current problems is the scientific substantiation of the degree of soil drought tolerance of medium-fibre cotton varieties and their protective adaptive properties in the soil and climatic conditions of the middle and lower regions of the Zarafshan oasis [8,9,10].

Research is being conducted around the world on the selection of varieties with high yield and quality and drought tolerance, including based on the following priorities: the creation of cotton varieties with beneficial economic characteristics, resistance to adverse environmental factors; increase in the adaptive capacity of cotton; development of scientifically based recommendations for planting in areas with water shortages with different soil and climatic conditions; identification of drought and salt-tolerant genotypes based on selection and molecular methods; development of methods to increase the resistance of plants through the use of biologically active substances and trace elements [11,12].

Bukhara-6, Aqdarya-6, Bukhara-8, S-6524 and Bukhara-102 varieties of medium fibre cotton were used as the object of research. The study aims to determine the degree and form of protective adaptation of cotton varieties to soil drought based on the physiological and biochemical properties of water exchange, as well as to develop methods to increase drought tolerance.

Research tasks are seed germination rate under different soil moisture conditions, amount of water forms in leaves, transpiration rate, leaf diffusion resistance, diurnal and residual water shortages, water potential and turgor resistance of leaves, determination of proline and carbohydrate content in leaves, the number of chlorophylls and the degree of their binding to protein-lipid compounds, photosynthesis and respiratory rate, determination of the number of free amino acids and phenolic compounds and the growth of cotton varieties of soil drought, determine the impact on development and productivity, as well as to study and apply the effect of the electrification method on the yield and its quality to increase the drought resistance of cotton.

MATERIALS AND METHODS

The soil of the experimental field belongs to the alluvial meadow type, the depth of groundwater is 2-3 meters. Based on the pre-irrigation soil moisture, bulk density, and moisture capacity, the degree of moisture deficiency in the soil was determined and irrigation standards were set. The experimental sites were divided into 3 sections. The experiments were performed in four repetitions. The experiments were carried out based on agro-techniques adopted on farms. Fertilizers were given during ploughing, along with planting, and during plant growth (3 times). The total amount of fertilizers applied per hectare is nitrogen-250, phosphorus-175, and potassium-100 kg. Phenological observations, calculations and research work on plant growth and development were carried out in accordance with the methods of the Uzbek Cotton Research Institute (UzCRI). Determination of all physiological parameters and phenological observations were carried out in the experiments during the stages of weeding, flowering and germination of cotton. A fourth leaf developed from the third part of the main stem was taken for the study.

Physiological and biochemical processes of drought tolerance of cotton varieties and indicators of protective adaptive properties were determined using methods generally accepted in plant physiology and biochemistry.

RESULTS AND DISCUSSION

During scientific studies, the diffusion resistance of cotton leaves under conditions of varying soil moisture levels was studied during the mowing, flowering, and budding stages of the plant. Diffusion resistance of leaves is an indicator of plant photosynthetic productivity, which is inextricably linked with transpiration intensity and is one of the protective adaptive reactions of plants to drought [13,14].

The diffusion resistance of the leaves increased in all variants of the cotton from the combing to the budding stage. As the soil moisture level decreased, the diffusion resistance of the leaves also became higher. It was observed that the value of diffusion resistance was higher in the

experimental variant with 30% soil moisture than in the other variants.

Leaf diffusion resistance increased by 27.2% in the experimental variant with 50% soil moisture and 57.1% in the experimental variant with soil moisture at 30% compared to the control stage. In the flowering stage, it was 13.2 and 38.6 per cent, respectively, and in the flowering stage, it was 20.3 and 39.8 per cent. Carbohydrate metabolism plays an important role in increasing the resistance of plants to drought and high temperatures. During the experiments, the amount of glucose, sucrose, maltose and starch in the flowering stage of the cotton plant was determined. Under moderate soil moisture conditions, a decrease in glucose content compared to other options was observed, and under the influence of drought, a sharp increase in glucose content was observed.

In the variant with 30% soil moisture, glucose levels were found to be 147.5% higher than the control. In the context of soil drought, sucrose was 140.2% higher than the control. In particular, in the variant with soil moisture of 30%, the value of maltose reached 284.6% compared to the control.

The decrease in soil moisture level also led to a decrease in the amount of starch. Soil moisture was higher in the moderate variant than in both experimental variants. The reduction in starch content in the variant with 50% soil moisture compared to the control was 68.3%, and in the variant, with 30% soil moisture was 49%. Taking into account the sum of the detected carbohydrates, it was noted that their content increased by 127.3% compared to the control in the conditions of soil drought. In the variants with soil moisture of 70 per cent, the total water content in all-cotton varieties, including free water, was also high, but the amount of bound water was low. At 30 per cent humidity, the opposite was true.

In all cotton varieties, a decrease in total and metabolic water content, and an increase in the amount of bound water, from mowing to weaning under two different humidity conditions were found. Under conditions of limited humidity (30 per cent), there are significant differences in the amount of bound water compared to varieties in moderate humidity (70 per cent). Differences were also observed in the cross-section of varieties in terms of the amount of water-bound. Especially Bukhara-6, Bukhara-102 varieties differ from other varieties due to the high amount of bound water.

In addition to the daytime water shortage in the leaves of cotton varieties, residual water deficiency was also studied during vegetation experiments. Looking at the data obtained on the amount of residual water, it was observed that its value depends on the level of soil moisture and the stages of development of varieties. It was found that the value of residual water deficiency was high in all varieties under conditions of limited soil moisture (30 per cent). It was noted that the value of this indicator varies across varieties. In the conditions of soil drought, the value of residual water shortage was the highest in the varieties Aqdarya-6 and S-

6524, and the lowest in the varieties Bukhara-6 and Bukhara-102. For invariants with moderate soil moisture levels, the value of this indicator decreased. It was noted that the concentration of cell sap of Bukhara-6, Bukhara-102 and Bukhara-8 varieties is higher than in other varieties under both humidity conditions. Such a feature may be one of the protective properties aimed at providing plants with more water even in adverse conditions. A decrease in soil moisture level affected an increase in osmotic pressure in the cells. In all varieties from the beginning to the end of the vegetation was observed a different increase in the value of this indicator.

The amount of chlorophyll and the degree of its binding to protein-lipid compounds were also studied, and it was noted that the total and bound chlorophyll content is higher in Bukhara-6, Bukhara-102 and Bukhara-8 varieties in soil drought conditions. Based on the results of laboratory, vegetative and field experiments, the laws of protective adaptation of cotton to drought at the cellular, tissue and ontogenetic levels were determined.

Physiological and biochemical comparative characteristics of drought adaptation have been developed based on the mechanisms of physiological adaptation of cotton to drought - reduction of water consumption, accumulation of low molecular weight osmoprotectants, changes in metabolism, increased water use efficiency.

Due to the positive effects of electrification on the physiological and biochemical and water exchange processes, it was found that due to the increase in water storage, reduction of diurnal and residual water shortages, drought tolerance of cotton, yield and its quality increased in soil moisture deficiencies. and a rapid method for determining diffusion resistance, bound water in leaves, and amount of bound chlorophyll were developed.

To obtain a high and high-quality harvest in cotton farms located in the middle and lower regions of the Zarafshan oasis (with arid soil), drought-resistant and high-yielding cotton varieties Bukhara-6, Bukhara-102 and Bukhara-8 have been introduced. A rapid method for determining the resistance of a cotton plant to soil water scarcity has been developed and recommended for use. The use of electro-ignition methods to increase the drought tolerance, yield and quality of cotton varieties has been introduced. The scientific significance of the results of the study serves to develop the physiological and biochemical basis of adaptation of cotton varieties to different types of cellular, tissue and ontogenetic levels of protective responses and adaptation to soil drought, depending on the degree of soil drought. Drought-resistant varieties Bukhara-6, Bukhara-102 and Bukhara-8 were planted in arid areas with high temperatures and water shortages.

CONCLUSION

The laws of protective adaptation of cotton to drought at the cellular, tissue and ontogenetic levels were determined,

physiological, biochemical, habitual forms of drought tolerance of varieties depending on soil moisture level were scientifically based and on this basis a model of drought-tolerant cotton varieties was created.

Physiological and biochemical comparative characteristics of drought adaptation have been developed based on the mechanisms of cotton adaptation to drought - reduction of water consumption, accumulation of low molecular weight osmoprotectants, changes in metabolism, and increased water use efficiency.

The reaction of cotton varieties to soil drought depends on the characteristics of the variety, the level of water supply to the physiological and biochemical processes of plants, as well as the impact on yield and quality.

All cotton varieties grown under conditions of moderate soil moisture were observed to have significantly lower daytime and residual water shortages, leaf water potential, the osmotic pressure of cell sap, protoplasm viscosity, cell dehydration, and heat resistance than plants grown under soil drought conditions. Soil drought has led to a relative increase in the amount of bound water in all-cotton varieties, water scarcity in the leaves, protoplasmic viscosity, and dehydration and heat resistance of leaf cells. It was noted that the value of physiological and biochemical indicators of drought tolerance is highest in varieties resistant to drought. A rapid method for determining the amount of residual water deficiency and diffusion resistance in the leaves, the amount of bound water in the leaves, and the amount of bound chlorophyll was developed and proposed to determine the degree of resistance of the cotton plant to soil water deficiency. In years of water scarcity, and atmospheric and soil drought, the use of environmentally friendly electrification methods has led to an increase in drought tolerance, yield and quality of cotton varieties.

It was recommended to sow Bukhara-6, Bukhara-102, Bukhara-8 varieties of cotton, which are drought-resistant, high yield and high quality, to obtain high and high-quality crops in cotton farms in drought-prone areas. High-quality and high-quality yields in Bukhara-6, Bukhara-102 and Bukhara-8 cotton varieties were determined in the order of 1-2-1, S-6524, Akdarya-6 varieties in the order of 1-3-1 irrigation, and cotton yield in Bukhara-6 variety were 38.4–40.7, Bukhara-102 variety 37.5–38.5 and Bukhara-8 variety 35.3–36.4 quintals, based on which the fibre quality meets international standards.

ACKNOWLEDGEMENTS

The authors acknowledge the immense help received from the scholars whose articles are cited and included in references to this manuscript. The authors are also grateful to the authors/editors/publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

We also express our deep gratitude to the leadership and scientific staff of the Uzbek Cotton Research Institute and

Bukhara State University for their close assistance in conducting the research.

The authors report no conflicts of interest.

Financing: The Source of funding is nil.

REFERENCES

- Khan A, Pan X, Najeeb U, Tan DK, Fahad S, Zahoor R, Luo H. Coping with drought: stress and adaptive mechanisms, and management through cultural and molecular alternatives in cotton as vital constituents for plant stress resilience and fitness. *Biological research*. 2018; 51.
- Mahmood T, Khalid S, Abdullah M, Ahmed Z, Shah MK, Ghafoor A, Du X. Insights into drought stress signalling in plants and the molecular genetic basis of cotton drought tolerance. *Cells*. 2019 Dec 31; 9(1): 105.
- Toshtemirovna N.U., Ergashovich K.A. Physiology, productivity and cotton plant adaptation under the conditions of soil salinity. *International Journal of Recent Technology and Engineering*. 2019; 8(2 S3): 1611-3.
- Norboyeva U.T., Kholliyev A.E. Salinification influence on physiology of water exchange in cotton plant varieties (*Gossypium Hirsutum*L.). 2017; 7(41): 16-18.
- Ergashovich KA, Azamatovna BZ, Toshtemirovna NU, Rakhimovna AK. Ecophysiological effects of water deficiency on cotton varieties. *Journal of Critical Reviews*. 2020; 7(9): 244-6.
- Norboyeva U.T., Kholliyev A.E. soil salinity and saline tolerance of the sorts of cotton. In *Mechanisms of resistance of plants and microorganisms to unfavourable environmental*. Irkutsk, July 10-15, 2018; (PART I): 567- 570.
- Magwanga RO, Lu P, Kirungu JN, Lu H, Wang X, Cai X, Zhou Z, Zhang Z, Salih H, Wang K, Liu F. Characterization of the late embryogenesis abundant (LEA) proteins family and their role in drought stress tolerance in upland cotton. *BMC genetics*. 2018 Dec; 19(1): 1-31.
- Norboyeva U.T., Kholliyev A.E. water interchange and saline tolerance of the sorts of cotton. In *Mechanisms of resistance of plants and microorganisms to unfavourable environmental*. Irkutsk, July 10-15, 2018; (PART I): 563-566.
- Ackerson RC. Osmoregulation in cotton in response to water stress: II. Leaf carbohydrate status in relation to osmotic adjustment. *Plant physiology*. 1981 Mar; 67(3): 489-93.
- Turner NC, Begg JE. Plant-water relations and adaptation to stress. *Plant and soil*. 1981 Feb; 58(1): 97-131.
- Toshtemirovna N.U., Ergashovich K.A. Regulation of the water balance of the cotton varieties under salting conditions. *ACADEMICIA: An International Multidisciplinary Research Journal*. 2019; 9(8): 5-9.
- Ergashovich KA, Toshtemirovna NU, Rakhimovna AK, Abdullayevna FF. Effects of microelements on drought resistance of the cotton plant. *International Journal of Psychosocial Rehabilitation*. 2020; 24(2): 643-8.
- Huang B. Role of root morphological and physiological characteristics in drought resistance of plants. *Plant-Environment Interactions*. Marcel Dekker Inc., New York. 2000 Jul 12: 39-64.
- Kholliyev A.E., Norboyeva U.T., Kholov Y.D., Boltayeva Z.A. Productivity of cotton varieties in soil salinity and water deficiency. *The American Journal of Applied sciences*. 2020; 2(10): 7-13.
- Yogesh Hole et al 2019 *J. Phys.: Conf. Ser.* 1362 012121.
- Zahoor R, Dong H, Abid M, Zhao W, Wang Y, Zhou Z. Potassium fertilizer improves drought stress alleviation potential in cotton by enhancing photosynthesis and carbohydrate metabolism. *Environmental and Experimental Botany*. 2017 May 1; 137: 73-83.



Сведения об источнике

NeuroQuantology

Годы охвата Scopus: с 2007 по настоящий момент

Издатель: Anka Publishers

ISSN: 1303-5150

Отрасль знаний: [Physics and Astronomy: Atomic and Molecular Physics, and Optics](#) [Neuroscience: Cognitive Neuroscience](#) [Neuroscience: Developmental Neuroscience](#)

Тип источника: Журнал

[Посмотреть все документы >](#)

[Настроить уведомление о документах](#)

[Сохранить в список источников](#) Source

Homepage

CiteScore 2021
1.3



SJR 2021
0.285



SNIP 2021
1.038



[CiteScore](#) [CiteScore рейтинг и тренды](#) [Содержание Scopus](#)

Улучшенная методика расчета CiteScore

Рейтинг CiteScore 2021 отражает количество цитирований в 2018-2021 гг. статей, обзоров, материалов конференций, глав книг и информационных документов, опубликованных в 2018-2021 гг., деленное на количество публикаций за 2018-2021 гг. [Подробнее >](#)

CiteScore 2021

$$1.3 = \frac{1\,044 \text{ цитирований за 2018 - 2021 гг.}}{781 \text{ документов за 2018 - 2021 гг.}}$$

Вычисление выполнено 05 May, 2022

CiteScoreTracker 2022

$$2.0 = \frac{821 \text{ цитирований на текущую дату}}{419 \text{ документов на текущую дату}}$$

Последнее обновление 05 October, 2022 • Обновляется ежемесячно

Рейтинг CiteScore 2021

Категория	Рейтинг	Процентиль
Physics and Astronomy Atomic and Molecular Physics, and Optics	#160/205	22-й
Neuroscience Cognitive Neuroscience	#85/107	21-й
Neuroscience Developmental	#34/37	9-й

[Просмотр методики CiteScore >](#) [Часто задаваемые вопросы о CiteScore >](#) [Добавить CiteScore на свой сайт](#)



NEW

Check out our new metric to help you evaluate journals!

Dismiss

Learn More

General Information

Web of Science Coverage

Journal Citation Report

Peer Review Information

PubMed® Information

Return to Search Results

JOURNAL OF PHARMACEUTICAL NEGATIVE RESULTS

Share This Journal

ISSN / eISSN 0976-9234 / 2229-7723

Publisher RESEARCHTRENTZ ACAD PUBL EDUCATION SERVICES, 240 Elm Street, 2nd & 3rd Floors, Somerville, United States, MA, 02144

General Information

Journal Website Visit Site

1st Year Published 2010

Frequency Annual

Issues Per Year 1

Country / Region INDIA

Primary Language English

Web of Science Coverage

Collection	Index	Category	Similar Journals
Core Collection	Emerging Sources Citation Index (ESCI)	Pharmacology & Pharmacy	Find 5 Similar Journals



Search a topic within this journal**Search****Journal Citation Report™ (JCR)****Journal Citation Reports™ 2022****Journal Citation Indicator (JCI)****NEW METRIC**

The Journal Citation Indicator is a measure of the average Category Normalized Citation Impact (CNCI) of citable items (articles & reviews) published by a journal over a recent three year period. It is used to help you evaluate journals based on other metrics besides the Journal Impact Factor (JIF).

2021**0.13****Category:**

Pharmacology & Pharmacy

2020**0.18**

Category:

Pharmacology & Pharmacy

[Learn About Journal Citation Indicator](#)**Peer Review Information**

Publons Partner ⓘ	No, and this journal does not explicitly endorse Publons
Claimed Reviews on Publons ⓘ	12
Public Reports on Publons ⓘ	No
Signed Reports on Publons ⓘ	No
Publons Transparent Peer Review Partner ⓘ	No
Publons User Endorsements ⓘ	3

Sign up for a free [Publons](#) account to track your publications, citation metrics, peer reviews, and editing work for this journal.

PubMed® Information

Indexed In ⓘ	PubMed®
NLM® ID ⓘ	101616264
Corporate Author ⓘ	Gale Group.

Courtesy of the U.S. National Library of Medicine

Editorial Disclaimer: As an independent organization, Clarivate does not become involved in and is not responsible for the editorial management of any journal or the business practices of any publisher. Publishers are accountable for their journal performance and compliance with ethical publishing standards. The views and opinions expressed



journal are those of the author(s) and do not necessarily reflect the views or opinions of Clarivate. Clarivate remains neutral in relation to territorial disputes, and allows journals, publishers, institutes and authors to specify their address and affiliation details including territory.

Criteria for selection of newly submitted titles and re-evaluation of existing titles in the Web of Science are determined by the Web of Science Editors in their sole discretion. If a publisher's editorial policy or business practices negatively impact the quality of a journal, or its role in the surrounding literature of the subject, the Web of Science Editors may decline to include the journal in any Clarivate product or service. The Web of Science Editors, in their sole discretion, may remove titles from coverage at any point if the titles fail to maintain our standard of quality, do not comply with ethical standards, or otherwise do not meet the criteria determined by the Web of Science Editors. If a journal is deselected or removed from coverage, the journal will cease to be indexed in the Web of Science from a date determined by the Web of Science Editors in their sole discretion – articles published after that date will not be indexed. The Web of Science Editors' decision on all matters relating to journal coverage will be final.

Clarivate.™ Accelerating innovation.

© 2022 Clarivate

[Copyright Notice](#)

[Terms of Use](#)

[Privacy Notice](#)

[Cookie Policy](#)

[Настройки файлов cookie](#)

[Help Center](#)

Follow us:

