

Effect of salinity on ecological and physiological characteristics of winter wheat varieties

Askar Kholliyev^{1,*}, Umida Norboyeva^{1,2}, and Dilfuza Teshaeva^{1,2}

¹Bukhara State University, Bukhara, Uzbekistan

²Bukhara State Pedagogical Institute, Bukhara City, Piridastgir, 2, 200100, Uzbekistan

Abstract. In connection with the influence of salt stress on winter wheat varieties in the soil and climatic conditions of the Bukhara oasis, it is of practical importance to determine the physiological foundations of the level of resistance and adaptation, as well as to determine resistance to stress and develop methods for increasing resistance. In studies to assess the effect of salt stress on wheat, physiological, biochemical, plasmolytic, morphological, biometric, statistical, comparative-analytical, phenological, and gasometric methods were used. In the ontogeny of winter wheat varieties, physiological methods have been developed for determining the degree of resistance to salt stress by determining the amount of bound water in the leaves (BWC), the level of cell turgor (TC), physiological methods for increasing resistance to salt stress by treating seeds before sowing. Key words: salt stress, soil salinity, winter wheat varieties, physiological parameters, adaptation, resistance, cell turgor, productivity.

1 Introduction

Studying the problem of soil salinity is important, since the need of the world's population for food is increasing every year. The main anthropogenic factors causing soil salinization are the intensification of agriculture, improper agrotechnical measures, including the use of high rates of artificial fertilizers, improper irrigation, and deforestation. About 20% of agricultural land is at risk of soil salinization. Rapid climate change causes long-term droughts affecting more and more regions of Europe [1-2].

Soil salinization has become one of the major environmental issues worldwide and is expected to worsen due to projected climate change. Arid and semi-arid agricultural areas are particularly sensitive to the impact of climate change on increased soil salinity [3-4].

More than 954 million hectares of the total land area in the world (more than 20% of agricultural land) are subject to salinization, especially in arid and semi-arid regions [5].

An increased content of sodium in the soil is one of the most common causes of abiotic stress in plants [6].

Globally observed stress factors have a serious negative impact on living organisms, including plants. As a result, indicators of plant productivity and yield, as well as its

* Corresponding author: askarkholliyev@gmail.com

quality, are reduced. The most important tasks are considered to be the intensification of scientific research aimed at reducing the negative impact of such stress factors and the development of measures to preserve the lost crop, the assessment and justification of the physiological aspects of the action of adverse stress factors [7].

Studying the mechanisms of resistance of plants to salt stress and revealing it is one of the urgent theoretical and scientific problems in the world, by deepening scientific research in this field, creating methods of using exogenous and endogenous substances to increase the resistance of wheat to stress factors, wide use of the existing gene pool of crops in genetic-selection research and great attention is paid to wide application to agricultural production.

The need for such studies is explained by the fact that to overcome the stress caused by plant salinity, activation of various physiological and biochemical mechanisms is required [8].

Grain crops are one of the important industrial crops that provide raw materials for various industries. Optimal factors aimed at preserving the crop are necessary when growing products at the level of demand. Improving the agro-ameliorative state of irrigated lands of our republic, improving ecological, physiological and agrotechnical measures to prevent soil salinization, identifying, creating and putting into practice crop varieties adapted to stress factors, physiological and biochemical characteristics of wheat varieties that express the level of resistance and productivity in stressful conditions, and adaptation of varieties, certain results have been achieved in the evaluation and scientific substantiation of reactions [9-12].

2 Materials and Methods

The aim of the study is to assess the physiological features of the influence of salt stress on the physiological and productive indicators of winter wheat varieties under conditions of Bukhara region, and to develop physiological methods to determine the resistance of wheat to this factor and to improve it.

The varieties Grom, Pervitsa, Starshina, Alexeyevich, Krasnodarskaya-99, Vassa, Asr and Antonina belonging to the group of winter wheat varieties were used as an object of research.

Laboratory experiments were conducted at the Laboratory of Ecological Physiology of Bukhara State University, and all field experiments were conducted at Mashhura-Munojot, Boboeva Munavvar land, and Islohat-98 farms, Jondor district.

The intensity of transpiration was determined by weighing on a VT-500 torsion balance (Russia). The total moisture content of leaves was determined by the thermostatic method, and the density of cell sap was determined at noon on a digital refractometer RUO Shui 2GHS-L (Japan). The daily and residual water deficit in the leaves was determined by the Shmatko method with some modifications. The amount of total chlorophylls in leaves was determined using a Minolta SPAD-502 chlorophyllometer (SPAD-Soil Plant Analysis Development) (Japan). The intensity of half-leaf photosynthesis and the intensity of respiration by calculating the amount of released SO₂, the level of turgorecence of leaf cells were determined using the Turgoromer device (TN-10-60TS) (Moldova). Leaf area expansion was measured by cross sections, and net photosynthetic efficiency was calculated using the thermostat method.

All observations, measurements and studies of plant growth and development were carried out in accordance with the method of surveying crop varieties. Each experiment was performed in three repetitions at the biological and three repetitions at the analytical level. Statistical data processing was carried out using MS Excel 2003 and Sigma Stat software applications.

3 Results and Discussion

During the research, the data were obtained on the rate of transpiration, total leaf moisture, water deficiency of leaves, the content of bound water in leaves, the level of water retention in leaves, and the level of leaf turgor of the cells, cell sap density, the amount of chlorophyll pigments, the rate of photosynthesis and respiration in the studied varieties under salt stress.

3.1 The rate of transpiration

Due to transpiration, not only the evaporation of water through the leaf, but also the adsorption of water and the movement of water and substances dissolved in it throughout the plant are provided. In the course of the research, the transpiration of released winter wheat varieties Grom, Pervitsa, Starshina, Alekseevich, Krasnodarskaya-99, Vassa, Asr and Antonina at the stages of tuberization, flowering and milk maturation was studied.

High transpiration was noted in wheat varieties Grom, Starshina, Krasnodarskaya-99. In the control variants, compared with the experimental ones, activation of water consumption by plant leaves was found. Compared with the stage of tuberization and flowering, the acceleration of water evaporation by plants was noted in all studied varieties in the phase of milky ripeness. It was also established that there are differences between varieties in terms of transpiration levels. It was noted that in varieties Starshina, Grom and Krasnodar-99 the intensity of transpiration was significantly higher than in varieties Antonina, Alekseevich and Vassa. The lowest results for this indicator were noted in the varieties Pervitsa and Asr. It was noted that the Starshina, Grom and Krasnodar-99 varieties evaporate more water than other varieties and activate their metabolic processes compared to other varieties.

3.2 Total water content in leaves

Determining the amount of water in plants, one can observe changes in the water balance of plants under the influence of unfavorable factors. Under the influence of soil salinity, an increase in total moisture content was revealed in all varieties. The amount of bound water is not the same in plants grown in saline environments. In particular, it was noted that the amount of bound water is maximum under conditions of medium-high salinity of the soil.

During the experiments, the amount of water in the leaves of the studied varieties varied depending on salinity. High values of total moisture content were noted in Grom, Starshina, Krasnodarskaya-99 varieties. According to these indicators, varieties Alekseevich, Vassa, Antonina took an intermediate place. Salt-tolerant varieties store a large amount of water in leaf cells, and under conditions of water deficit, this was manifested in all experiments.

3.3 The amount of bound water in the leaves

The amount of bound water in the leaves was determined at the stages of tuberization, flowering, and milk maturation in wheat varieties. According to the numerical data determined during the experiments and presented in the table, it was found that the amount of bound water in the leaves varies to varying degrees depending on the stage of growth and development of varieties and the concentration of salts in the soil.

It is noted that the value of this indicator increases from the tuber stage to the milking stage in all varieties. Differences between tubulation and flowering stages were minimal. In all varieties, soil salinity increased the value of bound water content. At the same time, the rate of such growth was high in varieties Krasnodar-99, Grom and Starshina.

Similar correlations were observed at other stages of cultivar development. The change in the value of this indicator in the context of varieties, stages of growth and development, and in the range of salt concentrations can be determined by the ratio of varieties to unfavorable abiotic factors, i.e. soil salinity.

3.4 Intensity of Photosynthesis

The study of the effect of soil salinity on the physiological and biochemical parameters of winter wheat varieties reveals the ability of these wheat varieties to adapt to salt stress. The intensity of photosynthesis of all wheat varieties was studied at the stages of tuberization, flowering, and milk maturation under two different conditions: non-saline and moderately saline soils.

According to the data obtained on the rate of photosynthesis of wheat varieties, it was noted that the value of this indicator varies depending on the concentration of salts in the soil, as well as the level of growth and development of wheat varieties.

In the course of our experiments, it was found that the rate of photosynthesis increased in all varieties of wheat grown in the control variant, and the rate of photosynthesis decreased in all varieties in the experimental variants. Differences between varieties and in the value of this indicator were revealed. A high rate of photosynthesis was noted in the varieties Starshina, Grom and Krasnodar-99 in the cut of varieties with medium-high salinity. The intensity of photosynthesis was significantly lower in varieties Asr and Pervitsa.

3.5 Respiration Intensity

In the course of our studies, along with indicators characterizing the salt-tolerant properties of wheat, we also studied the respiration rate. Because the activity of metabolic processes in plants is directly related to the intensity of this indicator.

In the experiments, we studied the intensity of respiration of wheat varieties in the phases of tuberization, flowering, and milky ripeness. The intensity of respiration in the control variants was low, and in the experimental variants of all varieties, its increase was noted. The intensity of respiration was the highest in the experimental variant of variety Asr and increased in all varieties from the phase of tuber maturation to milk. In all studied wheat varieties, it was found that the magnitude of the respiration intensity increases in the tuber phase compared to the control.

Based on the data obtained, it was noticed that the value of the respiratory rate directly depends on the level of soil salinity. The intensity of respiration was the highest in the phase of milky ripeness of wheat varieties, and its rate was relatively low in the phase of tuberization and flowering. If we compare the differences in the context of varieties, then the varieties Pervitsa and Asr were distinguished by a high rate of respiration.

3.6 Impact of salinity on crops

Salinization of irrigated lands sharply reduces their productivity, which leads to a decrease in the gross harvest of agricultural crops and, first of all, wheat.

In field experiments, we studied the effect of medium-high soil salinity on the productivity of wheat varieties.

Based on the data obtained, it was found that the productivity of the studied wheat varieties depends on the level of soil salinity. Especially as a result of moderate salinization of the soil, a decrease in the mass of the crop was noted in all variants of the experiment. The level of yield of the studied wheat varieties was the highest in the control variants

compared to other variants of the experiment; a decrease in the yield weight was observed under conditions of soil salinity. In all variants of the experiment, a decrease in the mass of the crop was noted in all varieties compared to the control.

In the control variant, the wheat yield of Starshina was 64.2; 52.3 in the variety Pervitsa; variety Antonina 56.0; 62.5 in Grom variety; 50.7 for variety Vek and 57.8 for variety Alekseevich; 60.6 in Krasnodar-99 variety; Variety Vassa amounted to 56.5 centners. It was noted that the yield level of all studied wheat varieties in the medium-high saline variant was reduced compared to the control variant. In particular, the yield weight of the experimental variant of Starshina wheat is 62.3; 48.1 in the variety Pervitsa; 53.1 in variety Antonina; Thunder grade 60.3; 46.2 for variety Vek and 54.6 for variety Alekseevich; 58.2 in the variety Krasnodar-99; Variety Vassa amounted to 53.8 centners.

It was noted that the yield of varieties Starshina, Grom and Krasnodar-99 was the highest in conditions of moderately saline soils compared to other studied varieties. Under these conditions, the lowest results in terms of yield weight were noted in wheat varieties Pervitsa and Asr. According to these indicators, varieties Antonina, Alekseevich and Vassa took an intermediate place.

During the experiments, it was noted that the values of physiological and productivity indicators determined in winter wheat varieties differ depending on soil salinity and biological characteristics of the varieties, and also the physiological aspects of the mechanism of the influence of salt stress on winter wheat varieties were analyzed. According to the above data, the first negative effect of salinity was manifested in the level of water exchange of plants.

Changes in the process of water exchange in winter wheat led to an increase in some physiological processes and a decrease or slowdown in others. At the same time, the water deficiency of the leaves, the amount of bound water in the leaves, the degree of water-retaining capacity of the leaves, the degree of cell turgor, the degree of cell sap density and the intensity of respiration are due to the increase. At the same time, there was a decrease in the rate of transpiration, total water in the leaves, the amount of chlorophyll pigments, and the rate of photosynthesis. Such changes in physiological processes led to a decrease in the overall productivity of the plant.

It has been established that the growth rate, leaf area, net productivity of photosynthesis and crop weight of varieties under salt stress conditions are the main factors determining productivity and are interconnected. The decrease in indicators that determine productivity, first of all, reduced the weight of the biological, and at the next stage, the economic crop.

During the study period, practical recommendations were developed based on the resistance characteristics of varieties to areas of the Bukhara oasis with different levels of salinity and areas with high salt stress. In the ontogeny of winter wheat varieties, physiological methods have been developed to determine the degree of resistance of varieties to salt stress by determining the amount of bound water in leaves (BWC), the level of cell turgor (TC), and increasing resistance to salt stress by treating seeds before sowing.

The scientific significance of the research results is explained by the fact that the protective responses and adaptability of wheat varieties to salt stress depend on the level of stress, and the development of physiological methods for determining resistance to salt stress and its increase. The practical significance of the research results is determined by the possibility of planting varieties Starshina, Krasnodarskaya-99 and Grom resistant to salt stress in areas with medium-high salinity, as well as varieties Antonina, Alekseevich, Vassa, which have medium resistance, in areas where the effect of stress factors is relatively weak, and the possibility of obtaining a quality crop. The implementation of the recommendations will improve the agricultural practices of growing winter wheat in areas with salinity.

4 Conclusion

The effect of soil salinity on the activity of physiological processes in wheat varieties was different depending on varietal characteristics. Compared to the control, the amount of total chlorophylls and the rate of photosynthesis decreased. At the same time, it was found that the intensity of respiration, the amount of bound water were higher under the influence of soil salinity compared to the control in all variants of the experiment. The yield of wheat varieties depends on the effect of salinity, and medium-high soil salinity adversely affected the morphological and physiological characteristics of varieties. In a saline environment, the growth of all varieties slows down and leaf surfaces become smaller. The rate of accumulation of dry matter in wheat was lower with soil salinity compared to the control. The net productivity of photosynthesis was estimated depending on soil salinity and biological characteristics of varieties. In salt-tolerant varieties Starshina, Grom and Krasnodar-99, the net productivity of photosynthesis and the decrease in crop weight under the influence of salinity were lower than in other studied varieties. The impact of soil salinity also caused a decrease in the biological and economic productivity of all studied wheat varieties. It is proved that the characteristics of resistance of varieties to soil salinity are different depending on their biological and individual characteristics. As a result of exposure to salt stress, the yield level of all varieties decreased. At the moment, varieties Starshina, Grom and Krasnodar-99 resistant to this factor did not show sharp changes in indicators characterizing salt tolerance and crop weight. In all experiments higher productivity of wheat varieties Starshina, Grom and Krasnodar-99 in comparison with other varieties in soil salinity conditions was revealed. Sharp decrease of productivity of varieties Pervitsa and Asr is connected with conditions of moderate and strong soil salinization. In variants treated before sowing wheat seeds with 3.5 % solutions of table salt, yield weight averaged 64.4 cwt. At the same time, compared to the control, the weight gain increased by 5.05%. The values of such indicators as bound water content in leaves (BWC), residual water deficit in leaves (RWD) and net photosynthetic productivity (NPP) were evaluated as higher compared with the control variant.

Physiological methods for determining the salt tolerance of wheat and increasing salt tolerance have been developed, and the positive effect of these methods on the productivity of wheat plants has been identified and proposed for production.

Based on the results obtained during the research, the following varieties of winter wheat can be recommended for sowing in medium and highly saline areas of the Bukhara region in terms of salinity resistance: -In moderately saline areas of the Bukhara region, it is recommended to plant varieties Starshina, Grom and Krasnodar-99 resistant to stress factors and salt, and in moderately saline areas - varieties Antonina, Alekseevich, Vassa.

Due to the low level of resistance to stress factors and soil salinity, winter wheat varieties Pervitsa and Asr are recommended to be planted on non-saline or slightly saline soil areas. use physiological methods to quickly determine the level of salt tolerance and increase resistance to salt stress.

References

1. R Munns, *Plant Cell Environ.* **25**. 239-250 (2002)
2. F Brini, I Amara, K Feki, M Hanin, H Khoudi, K Masmoudi, *Acta Physiol. Plant.* **31**. 145-154 (2009)
3. Okur B, Oren N *Soil salinization and climate change*// In *Climate Change and Soil Interactions*. Elsevier. -pp. 331-350 (2021)
4. D L Corwin, *Eur. J. Soil Sci.* **72(3)**. 842-862 (2021)

5. Munns R, Tester M //Annu.Rev.Plant Biol. **59** 651-681 (2008)
6. Shahid S, Zaman M, Heng L *Soil salinity: Historical perspectives and a world overview of the problem*// In Guideline for Salinity Assessment, Mitigation and Adaptation Using Nuclear and Related Techniques; Springer: Cham, Switzerland. -pp. 43-53 (2018)
7. Kholliyev A E, Teshaeva D R, Ra journal of applied research **08**. 03. 209-213 (2022)
8. Norboyeva U T, Kholliyev A E, International Journal of Recent Technology and Engineering (IJRTE) **8**, 2. S3, 1611– 1613 (2019)
9. Kholliyev A E, Norboyeva U T et.al., International Journal of Current Research and Review **13**. 04. 4-7 (2021)
10. Hamroyev A., Jumayeva H., E3S Web of Conferences **420**. 10007 (2023)
11. Kuldoshev R. et al., E3S Web of Conferences **371**. 05069 (2023)
12. Qo‘ldoshev A. R., Psychology and Education Journal **58**. 1. 4981-4988 (2021)