




BUXORO DAVLAT UNIVERSITETI ILMIY AXBOROTI

Научный вестник Бухарского государственного университета
Scientific reports of Bukhara State University

4/2022



E-ISSN 2181-1466



9 772181 146004

ISSN 2181-6875



9 772181 687004



4/2022

**PUBLISHED
SINCE 2000**
(Online since 2020)

**PUBLISHED SIX
TIMES A YEAR**

2022/4(92)

CHAIRMAN OF THE EDITORIAL BOARD:
Khamidov O.Kh.

Doctor of Economics, Professor

EDITOR-IN-CHIEF:

Zaripov G.T.

Candidate of technical sciences, Docent

INTERNATIONAL EDITORIAL BOARD:

Kuzmichev N.D. (Russia)

Doctor of Physics and Mathematics, Professor

Danova M. (Bulgaria)

Doctor of Philology, Professor

Margianti SE. (Indonesia)

Doctor of Economics, Professor

Wünsch Th. (Germany)

History of E.Europe Dr. of phil. habil, Professor

Minin V.V. (Russia)

Doctor of Chemical Sciences

Tashkaraev R.A. (Kazakhstan)

Doctor of Technical Sciences

Muminov M.E. (Malaysia)

Candidate of Physics and Mathematics

Srivastava P.K. (India)

American and English Literature PhD in English

NATIONAL EDITORIAL BOARD:

Adizov B.R.

Doctor of Pedagogical sciences, Professor
(Deputy Editor-in-Chief)

Abuzalova M.K.

Doctor of Philological sciences, Professor

Amonov M.R.

Doctor of Technical sciences, Professor

Barotov Sh.R.

Doctor of Psychological sciences, Professor

Bakoyeva M.K.

Doctor of Philological sciences

Buriyev S.B.

Doctor of biological sciences, professor

Djurayev D.R.

Doctor of Physics and Mathematics, Professor

Durdiyev D.K.

Doctor of Physics and Mathematics, Professor

Olimov Sh.Sh.

Doctor of Pedagogical sciences, Professor

Kakhkhorov S.K.

Doctor of Pedagogical sciences, Professor

Umarov B.B.

Doctor of Chemical sciences, Professor

Urayeva D.S.

Doctor of Philological sciences, Professor

Rashidov O.R.

Doctor of Historical sciences, Docent

DEPUTY EDITORS-IN-CHIEF:

Navruz-zoda B.N.

Doctor of Economics, Professor

Turayev H.H.

Doctor of Historical sciences, Professor

Juraev N.K.

Doctor of Political sciences, Professor

Jumaev R.G.

PhD in Political sciences, Docent

Kuvvatova D.Kh.

Doctor of Philological sciences, Professor

Akhmedova Sh. N.

Doctor of Philological sciences, Professor

**SCIENTIFIC REPORTS OF
BUKHARA STATE
UNIVERSITY**

**BUXORO DAVLAT
UNIVERSITETI ILMIY
AXBOROTI**

**НАУЧНЫЙ ВЕСТНИК
БУХАРСКОГО
ГОСУДАРСТВЕННОГО
УНИВЕРСИТЕТА**

*The journal is published in the Bukhara
Regional Department of Press and
Information of the Press and Information
Agency of Uzbekistan on August 24, 2020
With registered certificate № 1103*

*The journal "Scientific reports of Bukhara
state university" is included in the list of
scientific publications recommended to
publish the main scientific results of
doctoral dissertations of the Higher
Attestation Commission under the
Cabinet of Ministers of the Republic of
Uzbekistan on philology and physical and
mathematical sciences.*

*The journal is intended for professors
and teachers of higher educational
institutions, senior researchers, students,
scientific staff of scientific research
institutions, teachers of academic
lyceums, professional colleges, as well as
researchers in general secondary
education and various fields.*

**Founder: BUKHARA STATE
UNIVERSITY**

Executive secretary:

Sayfullaeva N.Z.

**Doctor of Philosophy in
Pedagogical Sciences (PhD)**

Editor: Sobirova Z.R.

Department technicians:

Shirinova M.Sh.

Raximova S.M.

CONTENTS

EXACT AND NATURAL SCIENCES		
Safarov J. Sh., Juraeva M. M.	Investigation of the hyperbolic type integro-differential equation in a square by the fourier method	2
Элмурадова Х. Б.	Обратная задача для псевдопараболических интегродифференциальных уравнений	10
Nazarova S.M.	Mechanical and microaggregate composition of irrigated meadow soils of the karakul oasis	24
Turdiyev Kh. Kh., Boltaev A. A.	The problem of determining the one-dimensional kernel of the viscoelasticity system	30
Ahadov A. A.	Importance of hydrogen energy and some methods of its production	54
LINGUISTICS		
Юлдашева Д. Н.	Сукутда ички нутқнинг воқеланиши	62
Рўзиев Я. Б.	Ўзбек тилшунослигида кўчирма ва ўзлаштирма нутқнинг талқину тавсифи	68
Jurayeva Z.R.	The image of a man and a woman in russian and uzbek paroemia with a gender component	77
Rasulov Z.I.	The role of the context and situation in the formation of elliptical structures	81
Собирова З. Р.	Диалектизм ҳодисаси ва унинг инглиз тили шеваларида кўлланилиши	87
Раджабова М. А.	Ономастик бирликларнинг фразеологизмлар таркибидаги гендер хусусиятлари	92
Turdiyeva Z.T.	The influence of gender on achievement in learning english as a second language	99
Kozieva I. K.	Actual and potential indicator anthroponyms in the russian language	104
Жураев Н. Дж.	Ўзбек ва инглиз тилларидаги “тил ва нутқ” мавзуй гуруҳидаги мақолларда синонимия, градуонимия, антонимия	112
Rahmatullayev Sh. N.	O‘zbek ertaklarida antroponimlarning lingvomadaniy asoslari	119
Ahmedov A. R., Po‘latova S. M.	Vobkent tumani shevasida so‘zning morfemik tarkibi	123
Barotova M. O.	About participation of paralinguistic tools with head	128
LITERARY CRITICISM		
Temirova Dj. Kh.	The genre of “Mockery” in russian and uzbek folklore	135
Худойкулова М. А.	Мустақиллик йилларида баҳсдаги адабий-назарий муаммолар талқини	146
Jamolova Z. N., Khudoykulova M. A.	The place of the literary review in the criticism of 20 th century	151
“NAVOIY GULSHANI”		
Sayliyeva Z.R.	Ideological-artistic analysis of the 10th verse in Alisher Navoi's “Badoye ul-bidoya” collection	155
HISTORY		
Raufov I.S.	Gpp Kandym - example cooperation between Uzbekistan and Russia in the oil and gas industry	159

MECHANICAL AND MICROAGGREGATE COMPOSITION OF IRRIGATED MEADOW SOILS OF THE KARAKUL OASIS

Nazarova S.M.

*Associate Professor of the Agronomy
and Soil Science Department,
Bukhara State University
nazarovasm@mail.ru*

Abstract. *The article shows the changes in the mechanical and microaggregate composition of the irrigated meadow soils of the Karakul oasis under the influence of anthropogenic factors. Morphological changes in the genetic layers of soils, in particular, the mechanical and microaggregate composition, are represented by methods adopted in soil science.*

The mechanical composition of soils has a significant impact on their following properties: physical, water, chemical, physico-mechanical, biological, thermal, etc. The moisture-retaining and load-bearing capacity of soils, heat transfer, physical and mechanical properties, soil resistivity during its processing, maturation time, viscosity, flexibility, subsidence, etc. directly depend on the mechanical composition. The mechanical composition of the soils in which scientific research is conducted is diverse, which is largely determined by the composition of the soil-forming parent rocks and human activity.

The mechanical and microaggregate composition of soils is the main indicator of morphological, reclamation, water-physical and physico-mechanical bonitation of soils and expresses the specifics for all genetic layers of soils of various types and types. According to the mechanical and microaggregate composition of soils, geomorphological areas with their own specifics are distinguished. These soils are mechanically composed of loam, light, medium, heavy loam.

Keywords: *anthropogenic factor, soil mechanical and microaggregate composition, medium, fine and large sand, large, medium and fine dust, particles*

Introduction. Extensive research work on soil fertility, its management and other properties has been carried out in foreign countries and in our republic. Irrigated meadow soils, common in the Karakul oasis, differ from each other in their textural features not only in soil areas, but also in soil and climatic conditions.

The morphogenetic structure, geographical location and reclamation condition, agrophysical and agrochemical properties of the soils of the Karakul oasis and other territories have been studied by a number of scientists of Uzbekistan. N. Feliciano [15], K. Gafurov, S. Abdullaev [1, 4], R. K. Kuziev [8, 9], L. A. Gafurova [5], N. M. Bobomurodov [3], R. Kurvantaev [6, 7, 10, 11], M. Umarov [13], H. T. Artikova [2], S. M. Nazarova [12], O. Scientific research was conducted by Sharipov [14]. However, scientific studies of the agrophysical properties of irrigated meadow soils of the Karakul oasis are currently insufficiently conducted.

Irrigated meadow soils common in the Karakul oasis, located in the lower reaches of the Zarafshan River, were chosen as **objects and methods of research**.

The research was carried out in soil-field and analytical laboratory conditions, using such manuals as "Theory and methods of soil physics", "Guidelines for conducting chemical and agrophysical analyses of soils during earth monitoring", "Methods of conducting field experiments", as well as the reliability of the data obtained is ensured using the Microsoft Excel program. A. Made on the basis of the Dospekhov manual "Field Opita technique".

The results of the study. The mechanical composition of soils has a significant impact on their following properties: physical, water, chemical, physico-mechanical, biological, thermal, etc. The moisture-retaining and load-bearing capacity of soils, heat transfer, physical and mechanical properties, soil resistivity during its processing, maturation time, viscosity, flexibility, subsidence, etc. directly depend on the mechanical composition. The mechanical composition of the soils in which scientific research is conducted is diverse, which is largely determined by the composition of the soil-forming parent rocks and human activity.

The mechanical composition of soils is the main morphological indicator, and for soils of all types, and therefore for all genetic layers that make up it, a specific mechanical composition is characteristic. For

EXACT AND NATURAL SCIENCES

example, sandy, sandy loam, sandy loam, (light, medium, heavy) and clay (light, heavy) mechanical compositions will be inherent in certain genetic layers and layers.

Of great practical importance is the mapping of meadow alluvial soils by mechanical composition, widespread and irrigated in the Karakul oasis. Meadow alluvial soils in cross section are not only sandy loam, sandy loam, loamy and clayey, but also have an extremely complex mechanical composition of layered-layered structure. As the top layer of the incision becomes heavier or lighter, there is a rapid change of layers. In such complex lithological conditions, a kind of nutrient, water-air and thermal order necessarily arises. In this regard, it is advisable not only to compile soil maps of irrigated lands in the oasis, but also to make maps on the mechanical composition of soils for each farm plot. This, in turn, makes it possible to solve on a scientific basis such a problem as improving the fertility of irrigated soils, especially their reclamation. It should be noted that the chemical composition of the soil, especially its absorption capacity, the amount of humus, etc., of course, strongly depends on the mechanical composition.

The data obtained indicate that according to the mechanical composition of soils, administrative districts and geomorphological areas with their own specifics are distinguished. The soils of the Karakul oasis belong to the soils formed in the lower reaches of the Zarafshan River and, according to their mechanical composition, are sandy loam, sandy loam, light, medium, heavy loam (Fig.).

Analysis of the data obtained shows that while medium and light loamy soils occupy the largest number of land plots (10509 ha), loamy and sandy loam soils of mechanical composition occupy 1177-1830 hectares. The soil sections of the Karakul oasis, located on the territory of the Lower Zarafshan, can be divided into 5 main groups according to their mechanical composition: uniformly light; uniformly heavy; layered, lightening from bottom to top; weighting soils of various mechanical composition from bottom to top. The change in the mechanical composition of soils in the cross section has a strong effect on the water-physical properties of soils and the salt regime.

Particles of different sizes in the soil vary widely in cross-section. The mechanical composition of irrigated meadow soils common in the Karakul oasis mainly consists of the following particles: coarse-grained sand (1-0.25 mm) up to 0.4-3.6%, medium sand (0.25-0.1 mm) 0.1-0.9%, fine sand (0.1-0.05 mm) 34.4-51.1%. Deposits, of which these soils consist, they also differ in a large content of fine grains of sand, the content of which in individual sections is 46.2-51.1%. The largest number of large dust particles (0.05-0.01 mm) accounts for 34.6-69.2%, the largest number of them scattered in farms is formed in layers with a clay mechanical composition, the smallest-on light soils. The cross-section between large dust particles (0.05-0.01 mm) is up to 32.2-35.6% higher than that of medium and fine dust. Il particles (less than 0.001 mm) make up to 2.7-9.6% and even in layers with heavy mechanical composition make up an insignificant amount. 2010 compared to 2017

from year to year, there is an increase in fine sand particles and a decrease in large dust particles. This is because the dust indicates that its particles were washed out under the influence of irrigation.

The mechanical composition of soils is characterized by:

- a) the complete absence of particles larger than 1 mm with a rough skeleton (pebbles) ;
- b) excess of the number of particles 0.1-0.01 mm;
- c) excess of large dust (0.05-0.01 mm) particles on most soils, the number of which in some cases can reach 35-69 % ;
- g) very low (0.1-3.6%) content of silt particles in loamy soils.

The above data indicate that it is clear from the old ones that the mechanical composition of irrigated soils weighs a little. This, of course, occurred as a result of the impact of anthropogenic factors on the surface layers of the soil. Most of the soils common in the farms of the area adjacent to the territory of sandy deserts belong to loamy and sandy loam of mechanical composition, the state of their civilization is somewhat lower.

The number of fine grains of sand (0.1-0.05 mm), in the Dargaly massif of the Karakul district, the above number of particles is distributed unevenly across the cross section. Its number ranges from 7.8 to 41.1% in 2010 to 34.4-51.1% by 2017. It should be noted that the number of particles in the soils listed above is distributed to varying degrees. For example, large dust particles (0.05-0.01 mm) make up 34-69%.

Irrigated soils with a variety of mechanical composition are also common in the Karakul oasis. Due to the fact that irrigated soils in the farms of the Karakul oasis are diverse in mechanical composition, agrotechnical (plowing, irrigation, tillage, fertilization, etc.), reclamation (leaching, installation of a system of collector ditches, registration of the depth of groundwater, etc.) measures carried out must be differentiated by mechanical composition.

Table 1. Mechanical and microaggregate composition of irrigated meadow soils of the Karakul oasis

Cuts №	Layer depth, cm	Fraction size, mm yes, quantity in %										
		>0,25	0,25-0,1	0,1-0,05	0,05-0,01	0,01-0,005	0,005-0,001	<0,001	Physical clay (<0,01mm)	Current number of units	Dispersion coefficient	
In 2010												
8	0-46	0,8	0,2	36,7	39,4	7,8	7,9	7,2	22,9	6,7	41,6	
		1,2	0,3	41,3	41,0	6,8	6,4	3,0				
		-0,4	-0,1	-4,6	-1,6	1,0	1,5	4,2				
	46-66	0,7	0,3	13,9	41,8	14,8	20,7	7,8	43,3	23,8	15,3	
		1,2	0,3	26,2	48,8	18,8	3,5	1,2				
		-0,5	0	-12,3	-7,0	-4,0	17,2	6,6				
	66-85	0,4	0,1	41,1	34,6	3,6	13,1	7,1	23,8	9,3	40,8	
		1,0	0,3	38,9	43,1	3,6	10,2	2,9				
		-0,6	-0,2	2,2	-8,5	0	2,9	4,2				
	85-106	0,8	0,2	17,7	69,2	5,1	4,3	2,7	12,1	7,8	66,6	
		0,4	0,1	12,5	77,0	5,1	3,1	1,8				
		0,4	0,1	5,2	-7,8	0	1,2	0,9				
	106-137	0,4	0,1	7,8	60,0	8,0	18,0	5,7	31,7	22,9	14,0	
		0,8	0,2	6,6	76,4	14,0	1,2	0,8				
		-0,4	-0,1	1,2	-16,4	-6,0	16,8	4,9				
	HCP- 613,8 P, %-0,01											
	In 2017											
	9	0-35	3,6	0,9	46,2	12,3	18,1	13,7	5,2	37,0	12,5	76,9
0,8			0,2	48,4	22,6	12,4	11,6	4,0				
2,8			0,7	-2,2	-10,3	5,7	2,1	1,2				
35-62		0,4	0,1	43,7	26,2	10,6	9,4	9,6	29,6	11,4	43,7	
		0,4	0,1	42,3	37,6	6,8	8,6	4,2				
		0	0	-1,4	11,4	-3,8	-0,8	-5,4				
62-90		0,8	0,2	37,5	32,2	6,9	18,0	4,4	29,3	10,6	63,6	
		0,5	0,1	42,4	37,9	6,7	9,6	2,8				
		0,3	0,1	-4,9	-5,7	0,2	8,4	1,6				
90-115		0,8	0,2	34,4	25,9	9,1	23,9	5,7	38,7	22,5	59,6	
		1,2	0,3	44,9	37,4	8,2	4,6	3,4				
		-0,4	-0,1	-10,5	-11,5	0,9	19,3	2,3				
115-147		0,8	0,2	43,0	35,6	6,3	11,0	3,1	20,4	11,7	70,9	
		0,4	0,1	54,7	31,2	5,3	6,1	2,2				
		0,4	0,1	-11,7	4,4	1,0	4,9	0,9				
147-186		0,8	0,2	51,0	29,7	8,3	6,6	3,4	18,3	16,5	70,5	
		1,2	0,3	40,6	39,1	5,2	11,2	2,4				
		-0,4	-0,1	10,4	-9,4	3,1	-4,6	1,0				
HCP- 423,9 P, %-0,01												

**Appendix: the figure of the table shows the mechanical composition of soils, in the denominator-the composition of the microaggregate.*

Study of the causes of natural soil fertility of the lower reaches of the Zarafshan River. Kurvantaev, 6, 7 showed that the high content of dust particles in these soils with a size of 0.05-0.01 mm and large stable microstructural elements with a size of more than 0.01 mm provides the best capillary pore formation, high moisture capacity and high water permeability.

Meadow soils consist mainly of fine grains of sand (0.1-0.05) and large dust particles (0.05-0.01 mm), microaggregate particles, so it is microstructural, which is characterized by the mobility of high moisture capacity and nutrients that provide the soil with good capillary porosity, a high amount of moisture, which in these cases determines high soil fertility.

When assessing from an agronomic point of view, it is sufficient to know not only the size of the mechanical elements of the soil, but also the presence of micro aggregates formed from some mechanical elements in the soil is characteristic. At the same time, it is important to study the comparative ability of these mechanical elements to wash under the influence of water. The description of sandy and clay soils is one of the important aspects in assessing soil structure, especially water permeability. The global aggregate composition of the soils of the Karakul oasis is diverse, which is associated with their genesis, morphology, soil-forming maternal sex and cultural irrigation activities of people. [12, 11, 14, 8].

If as a result of numerous studies, a large amount of factual data has been collected and the reasons for the formation of macroculture on irrigated soils have been studied through the use of crop rotation, siderates, additional organic and structure-forming substances. Irrigated soils of Uzbekistan are in a state of structure with alfalfa and other perennial herbs. Herbs have been found to have a positive effect.

It was noticed that the information we received, together with the full confirmation of the opinions of the above-mentioned scientists, was significantly divided into some aggregates as a result of irrigation and processing. Coarse sand (1-0.25 mm) 0.4-1.2%, medium sand (0.25-0.1 mm) 0.1 - 0.3%, fine sand (0.1 - 0.05 mm) 6.6 - 41.3% in 2010 and its amount increased to 40.6-54.7% in 2017. The sediments consisting of these soils are distinguished by the abundance of fine sand particles in the stream. The largest number of large dust particles (0.05-0.01 mm) was 41.0-76.4% in 2010, and by 2017 it was observed that their volume decreased by 22.6-39.1%. It can be seen that the amount of fine dust and silt granules in soils increased by 2017. For example, in the soils of the Dargali massif of the Karakul district in 2010, the content of fine dust was 1.2-10.2% and silt 0.8-3.0%, in 2017, respectively, 2.2-4.2%. As these results show, in the areas of Karakol, an increase in the number of real aggregates can be observed, that is, the process of structuring in the soil.

In short, the mechanical composition of the irrigated soils of the Spruce Oasis is somewhat lighter, the number of microaggregates is slightly smaller, and the decomposition coefficient is high.

The change in the surface layer of irrigated soils depends primarily on irrigation deposits and the composition of these deposits, which were formed as a result of irrigation through the Zarafshan River and the Amu-Karakul channels.

As a result of irrigation with turbid waters and other factors of human activity, a layer of agroirrigation was formed in a short time. It differs from the lower layers not only by its chemical properties, but also by its physical properties.

The mechanical and microaggregate composition of the Spruce Oasis is a key factor in the management of all processes in the soil, which, in turn, serves as a key indicator in the development of necessary measures for the effective use of soils. When processing, watering, fertilizing and placing various crops in the soil, it is necessary to stratify them taking into account their mechanical composition. From these results, the following conclusion can be drawn:

Conclusion. The mechanical and microaggregate composition of soils is the main indicator of morphological, reclamation, water-physical and physico-mechanical bonitation of soils and expresses the specifics for all genetic layers of soils of various types and types. According to the mechanical and microaggregate composition of soils, geomorphological areas with their own specifics are distinguished. These soils are mechanically composed of loam, light, medium, heavy loam.

The mechanical and microaggregate composition of irrigated meadow soils is dominated by the following mechanical elements: fine grains of sand (0.1-0.05 mm), coarse-grained (0.05-0.01 mm) particles, which in the section and geomorphological areas constitute a higher content compared to coarse, medium sand and medium, fine dust. In the farms of the district, where irrigated agriculture is still conducted, one can see some severity of the mechanical composition of distributed soils. This, of course, occurred as a result of the impact of anthropogenic factors on the surface layers of the soil. Most of the soils common in the farms of the area adjacent to the territory of sandy deserts belong to loamy and sandy mechanical composition, the state of their civilization is somewhat lower.

The mechanical and microaggregate composition of soils is as follows: incomplete content of large particles from 1 mm in a rough skeleton (rocky); a large number of particles 0.1-0.01 mm; excessive content of large dust (0.05-0.01) particles in most soils, the content of which in some cases reaches 35-50%, very small (2.7-6.6%) the number of fine particles. %).

References:

1. Абдуллаев С.А. *Агрофизическая свойства и солевой режим орошаемых почв оазисов Бухарской области. Автореф. канд. дисс.-Ташкент. 1975.– 34 с.*
2. Артикова Х.Т. *Бухоро воҳаси сугориладиган ўтлоқи аллювиал тупроқларининг умумий физик ва айрим сув-физик хоссалари, уларнинг аҳамияти. / Журнал “ЎЗМУ хабарлари” - №3/1. 2018.- Б. 47-51.*
3. Бобомуродов Ш.М. *Плодородие орошаемых почв нижнего течения реки Зарафшан и пути его повышения. Автореф. канд. дисс.- Ташкент. 2001.-25 с.*
4. Гафуров К., Абдуллаев С. *Характеристика почвенного покрова орошаемой зоны Бухарской области.- Ташкент: «Фан», 1982.- 130 с.*
5. Гафурова Л.А. *Изменение климата и проблемы сохранения и воспроизводства плодородия почв Узбекистана: реалии, тенденции и перспективы // Почвоведении в России: вызовы современности, основные направления развития: Материалы Всероссийской научно-практической конференции. – Россия, 2012,– С.104-109*
6. Курвантаев Р. *Плотность сложения почвы и урожайность хлопчатника. – Ташкент: «Узинформагрупп», -1991, - 88 с.*
7. Курвантаев Р. *Оптимизация и регулирование агрофизического состояния орошаемых почв пустынной зоны Узбекистана. Автореф. докт. дисс. -Ташкент. 2000. - С. 40-55.*
8. Қўзиев Р.Қ., Бобомуродов Ш.М. *Зарафшон қўйи оқими сугориладиган тупроқлари ва уларнинг унумдорлигини ошириши йўллари. –Тошкент: “Фан “. 2004.- 120 б.*
9. Кузиев Р.К., Сектименко В.Е. *Почвы Узбекистана.-Ташкент. 2009.-350 с.*
10. Курвантаев Р., Назарова С.М. *Агрофизическая характеристика орошаемых луговых почв Бухарского оазиса / Современные тенденции в научном обеспечении агропромышленного комплекса: Коллективная монография. [редкол.:Л.И. Ильин и др.; отв. за вып. В.В.Огорков]. - Иваново, 2019. - С. 91-95.*
11. Курвантаев Р., Назарова С.М. *Зарафшон воҳаси қўйи оқими сугориладиган ўтлоқи тупроқларининг агрофизикавий ҳолати. – Бухоро. 2021. - 126 б.*
12. Назарова С.М. *Бухоро воҳаси сугориладиган ўтлоқи тупроқларининг ҳозирги давр агрофизикавий ҳолати. Қишлоқ хўжалиги фанлари бўйича фалсафа доктори (PhD) автореферати.- Тошкент, 2019. - 44 б.*
13. Умаров М.У., Курвантаев Р. *Повышение плодородия орошаемых почв путём регулирования их физических свойств.- Ташкент: «ФАН», 1987.-106 с.*
14. Шарипов О.Б. *Бухоро вилояти иқлим шароитида агробиотехнологиялар асосида тупроқ унумдорлигини ошириши. / ҚарДУ хабарлари. №5.(37). - Қарши. 2018. - Б.120-122.*
15. Фелициант И.Н., Конобеева Г.М., Горбунов Б.В., Абдуллаев М.А. *Почвы Узбекистана. Бухарская и Навоийская области. -Ташкент: «Фан», 1984.- 153 с.*