

scitation.org/journal/apc

Volume 2365

**International Uzbekistan-  
Malaysia Conference on  
“COMPUTATIONAL MODELS AND  
TECHNOLOGIES (CMT2020)”  
CMT2020**

Tashkent, Uzbekistan • 24–25 August 2020

**Editors** • Rakhmatillo D. Alov, Kholmat M. Shadimetov,  
Abdullo R. Hayotov and Mirzoali U. Khudoyberganov



**AIP Conference Proceedings**

**AIP**  
Publishing

# AIP Conference Proceedings

---

Volume 2365

ISBN: 978-0-7354-4110-1

ISSN: 0094-243X

[scitation.org/journal/apc](http://scitation.org/journal/apc)

Proof

**AIP**  
Publishing

# International Uzbekistan- Malaysia Conference on “COMPUTATIONAL MODELS AND TECHNOLOGIES (CMT2020)” CMT2020

**Tashkent, Uzbekistan**  
24-25 August 2020

## Editors

**Rakhmatillo D. Aloyev**  
National University of Uzbekistan, Tashkent, Uzbekistan

**Kholmat M. Shadimetov**

**Abdullo R. Hayotov**  
V. I. Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

**Mirzoali U. Khudayberganov**  
National University of Uzbekistan, Tashkent, Uzbekistan

## Sponsoring Organizations

National University of Uzbekistan  
University Teknologi MARA (UiTM)  
V. I. Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences  
Navoi State Pedagogical Institute  
Samarkand Branch of Tashkent University of Information Technologies

All papers have been peer reviewed.



Melville, New York, 2021  
AIP Conference Proceedings

Volume 2365

To learn more about AIP Conference Proceedings visit <http://proceedings.aip.org>

## Editors

### Rakhmatillo D. Alov

National University of Uzbekistan  
Department of Computational Mathematics and Information Systems  
4, University Street  
Tashkent, 100174  
Uzbekistan

Email: alovr@mail.ru

### Mirzoali U. Khudoyberganov

National University of Uzbekistan  
Department of Computational Mathematics and Information Systems  
4, University Street  
Tashkent, 100174  
Uzbekistan

Email: mirzoali@mail.ru

### Kholmat M. Shadimetov

V.I. Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences  
Laboratory of Computational Mathematics  
4b, University Street  
Tashkent, 100174  
Uzbekistan

Tashkent State Transport University  
Informatics and Computer Graphics  
1, Odlikhodjaev Street  
Tashkent 100167  
Uzbekistan

Email: kholmatshadimetov@mail.ru

### Abdullo R. Hayotov

V.I. Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences  
Laboratory of Computational Mathematics  
4b, University Street  
Tashkent, 100174  
Uzbekistan

National University of Uzbekistan  
Department of Computational Mathematics and Information Systems  
4, University Street  
Tashkent 100174  
Uzbekistan

Email: hayotov@mail.ru

Authorization to photocopy items for internal or personal use, beyond the free copying permitted under the 1978 U.S. Copyright Law (see statement below), is granted by the AIP Publishing LLC for users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of \$30.00 per copy is paid directly to CCC, 222 Rosewood Drive, Danvers, MA 01923, USA: <http://www.copyright.com>. For those organizations that have been granted a photocopy license by CCC, a separate system of payment has been arranged. The fee code for users of the Transactional Reporting Services is: 978-0-7354-4110-1/21/\$30.00



© 2021 AIP Publishing LLC

No claim is made to original U.S. Government works.

Permission is granted to quote from the AIP Conference Proceedings with the customary acknowledgment of the source. Reproduction of an article or portions thereof (e.g., extensive excerpts, figures, tables, etc.) in original form or in translation, as well as other types of reuse (e.g., in course packs) require formal permission from AIP Publishing and may be subject to fees. As a courtesy, the author of the original proceedings article should be informed of any request for republication/reuse. Permission may be obtained online using RightsLink. Locate the article online at <http://proceedings.aip.org>, then simply click on the RightsLink icon/"Permissions/Reprints" link found in the article abstract. You may also address requests to: AIP Publishing Office of Rights and Permissions, 1305 Walt Whitman Road, Suite 300, Melville, NY 11747-4300, USA; Fax: 516-576-2450; Tel.: 516-576-2268; E-mail: [rights@aip.org](mailto:rights@aip.org).

ISBN 978 0 7354 4110 1  
ISSN 0004 243X  
Printed in the United States of America

*AIP Conference Proceedings, Volume 2365*  
International Uzbekistan-Malaysia Conference  
on “COMPUTATIONAL MODELS AND TECHNOLOGIES (CMT2020)”  
CMT2020

**Table of Contents**

<b>Preface: International Uzbekistan-Malaysia Conference on “Computational Models and Technologies (CMT 2020)”</b>	010001
<b>COMPUTATIONAL MATHEMATICS</b>	
<b>Lyapunov numerical stability of a hyperbolic system of linear balance laws with inhomogeneous coefficients</b> Rakhmatillo Alov and Dilfuza Nematova	020001
<b>Three-dimensional linear hyperbolic system</b> Rakhmatillo Alov and Dilfuza Nematova	020002
<b>Difference upwind scheme for the numerical calculation of stable solutions for a linear hyperbolic system</b> Nematova Dilfuza	020003
<b>An enhanced ant colony optimization with Gini index for predicting type 2 diabetes</b> Nur Hadirah Khairul Anwar, Rizauddin Saian, and Sumarni Abu Bakar	020004
<b>Group coloring via its geometric structures</b> Muhammed Bello and Nor Muhainiah Mohd Ali	020005
<b>Numerical treatment of nonlinear mixed Volterra-Fredholm integro-differential equations of fractional order</b> Zainidin Eshkuvatov, Zaid Laadjal, and Shahrina Ismail	020006
<b>Application of the Adams-Bashfort-Moulton method to the numerical study of linear fractional oscillators models</b> Parovik Roman	020007
<b>Effective finite-difference method for elastoplastic boundary value problems</b> Abduvali Khaldjigitov, Umidjon Djumayozov, and Sherzod Ibodullayev	020008
<b>Comparison of some numerical methods solutions of wave equations with strong dispersion</b> Dauletbay Utebaev and Bakhadir Utebaev	020009
<b>Numerical approach for nonlinear system of Fredholm-Volterra integral equations</b> Zainidin Eshkuvatov, Husnida Mamatova, Shahrina Ismail, Ilyani Abdullah, and Rakhmatillo Alov	020010
<b>Research of the hereditary dynamic Riccati system with modification fractional differential operator of Gerasimov-Caputo</b> Dmitriy Tverdyi and Roman Parovik	020011



<b>New iterative method for solving chemistry problem</b> Indranil Ghosh, M. S. H. Chowdhury, Suazlan Bin Mt Aznam, and Shukranul Mawa	020012
<b>Quasi random sequences for calculation multidimensional integrals and parallel algorithms</b> Abdujabar Rasulov, Matyokub Bakoev, and Makhmadrasul Rakhmatov	020013
<b>Weight optimal order of convergence cubature formulas in Sobolev space</b> Ozodjon Jalolov	020014
<b>The algorithm for constructing a differential operator of 2nd order and finding a fundamental solution</b> Ikrom I. Jalolov	020015
<b>On a Volterra type fractional integro-differential equation with degenerate kernel</b> Tursun K. Yuldashev	020016
<b>Monte Carlo methods for pricing weather derivatives</b> Abdujabar Rasulov and Gulnora Raimova	020017
<b>The white rat of numerical reproducibility</b> Michael Mascagni	020018
<b>Construction of cubature formulas with minimal number of nodes</b> Gaybulla Ismatullaev, Sayfiddin Bakhromov, and Ravshan Mirzakabilov	020019
<b>Optimal quadrature formulas with derivative for Hadamard type singular integrals</b> Dilshod Akhmedov and Kholmat Shadimetov	020020
<b>Optimal quadrature formulas for computing of Fourier integrals in <math>W_2^{(m, n-1)}</math> space</b> Abdullo Hayotov and Samandar Babaev	020021
<b>Optimal quadrature formulas with the trigonometric weight in the Sobolev space</b> Abdullo Hayotov and Bakhromjon Bozarov	020022
<b>On an exponential-trigonometric natural interpolation spline</b> Aziz Boltaev and Dilshod Akhmedov	020023
<b>An optimal quadrature formula in <math>K_2(P_3)</math> space</b> Siroj Azamov	020024
<b>Composite optimal formulas for approximate integration of weight integrals</b> Kholmat Shadimetov and Bahtiyor Daliyev	020025
<b>Numerical solution of Saint-Venant equations</b> Aziza Akbarova	020026
<b>An adequate computational model for a mixed problem for the wave equation in a domain with an angle</b> Mirzoali Khudoyberganov	020027

# Weight Optimal Order of Convergence Cubature Formulas in Sobolev Space

Ozodjon Jalolov<sup>a)</sup>

Bukhara State University, Bukhara 200117, Uzbekistan

<sup>a)</sup>Corresponding author: o.jalolov@mail.ru

**Abstract.** In this paper we investigate weight cubature formula in function spaces of S.L. Sobolev  $L_2^{(m)}, L_p^{(m)}, \tilde{L}_2^{(m)}$  for the functions defined in the  $n$ -dimensional unit cube  $K_n$  and obtain an upper estimate for the norm of error functionals of weight cubature formulas. The basis of theorem N.S. Bahvalov it is proved that considered weight cubature formulas are optimal on order of convergence in these spaces.

## INTRODUCTION

In many research papers examined the properties of optimal approximations of linear functionals [1-28], and others. In these papers the problem of optimality with respect to a certain space are investigated. Most of them are discussed in the Sobolev space [1]. Consider the cubature formula of the form

$$\int_{K_n} p(x)f(x)dx \approx \sum_{\lambda=1}^N C_\lambda f(x^{(\lambda)}), \quad (1)$$

in the space  $L_2^{(m)}(K_n)$ , where  $K_n$  is a  $n$ -dimensional unit cube. A generalized function

$$\ell_N(v) = p(v)\varepsilon_{K_n}(v) - \sum_{\lambda=1}^N C_\lambda \delta(v - v^{(\lambda)}), \quad (2)$$

is called a error functional of the cubature formula (1),

$$\langle \ell_N, f \rangle = \int_{K_n} p(x)f(x)dx - \sum_{\lambda=1}^N C_\lambda f(x^{(\lambda)}), \quad (3)$$

is an error of the cubature formula (1),  $p(x)$  is a weight function,  $\varepsilon_{K_n}(x)$  is characteristic function of  $K_n$ ,  $C_\lambda$  and  $x^{(\lambda)}$  are coefficients and nodes of the cubature formula (1),  $\delta(x)$  is the Dirac delta-function.

**Definition 1.** The space  $L_2^{(m)}(K_n)$  is defined as the space of functions, given on the  $n$ -dimensional unit cube  $K_n$  and having all the generalized derivatives of order  $m$ , square summable in norm [1]:

$$\|f/L_2^{(m)}(K_n)\| = \left\{ \int_{K_n} \sum_{|\alpha|=m} \frac{m!}{\alpha!} [D^\alpha f]^2 dx \right\}^{\frac{1}{2}}, \quad (4)$$

with the inner product

$$(f, \varphi)_{L_2^{(m)}(K_n)} = \int_{K_n} \sum_{|\alpha|=m} \frac{m!}{\alpha!} D^\alpha f \cdot D^\alpha \varphi dx,$$

where  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ ,  $dx = dx_1 dx_2 \dots dx_n$  and  $\alpha! = \alpha_1! \alpha_2! \dots \alpha_n!$ .

**Definition 2.** The cubature formula of the form (1) is called asymptotically optimal, if for the norms of error functional the following equality holds

$$\lim_{N \rightarrow \infty} \frac{\|\ell_N^{a, \sigma} / L_2^{(m)*}(K_n)\|}{\|\ell_N^a / L_2^{(m)*}(K_n)\|} = 1. \quad (5)$$

International Uzbekistan-Malaysia Conference on "COMPUTATIONAL MODELS AND TECHNOLOGIES (CMT2020)"

AIP Conf. Proc. 2365, 020014-1-020014-14; <https://doi.org/10.1063/5.0057015>

Published by AIP Publishing, 978-0-7354-4110-1/530.00

Here,  $\ell_N^o(x)$  and  $\ell_N^{o,o}(x)$  are error functional of optimal and asymptotically optimal cubature formulas of the form (1), respectively.

**Definition 3.** The cubature formula of the form (1) is called an optimal order of convergence, if for the norm of its error functional the following holds

$$\lim_{N \rightarrow \infty} \frac{\left\| \ell_N^{o,o} / L_2^{(m)^*}(K_n) \right\|}{\left\| \ell_N^o / L_2^{(m)^*}(K_n) \right\|} < \infty. \quad (6)$$

Here  $\ell_N^{o,o}$  is error functional of optimal order of convergence cubature formulas (1). In this paper we consider the problem of the descending order of norm of the error functional  $\left\| \ell_N / L_2^{(m)^*}(K_n) \right\|$ ,  $\left\| \ell_N / L_p^{(m)^*}(K_n) \right\|$  and  $\left\| \ell_N / L_2^{(m)^*}(K_n) \right\|$  with an increase in the number of its nodes. The results, which we obtain here, are to the arbitrary distribution of points.

### OPTIMAL IN ORDER OF CONVERGENCE OF WEIGHT CUBATURE FORMULAS IN THE SPACE $L_2^{(m)}(K_n)$

Here we explore weight cubature formulas, which are optimal for the convergence order. We have the following Here we explore weight cubature formulas, which are optimal for the convergence order. We have the following

**Lemma 1.** If for error functional (2) of cubature formula (1) satisfies the conditions

$$\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2) \cdot \dots \cdot \ell_{N_n}(x_n) \quad (7)$$

and

$$\left\| \ell_{N_i} / L_2^{(m_i)^*}(0,1) \right\| \leq c_i \frac{1}{N_i^{m_i}}, \quad (i = \overline{1, n}), c_i - constants \quad (8)$$

that is

$$\left\| \ell_N / L_2^{(m)^*}(0,1) \right\| \leq c_i O(h_i^{m_i}), \quad (i = \overline{1, n}), h_i = \frac{1}{N_i} \quad (9)$$

then

$$\left\| \ell_N / L_2^{(m)^*}(K_n) \right\| \leq c \cdot \frac{1}{\prod_{i=1}^n N_i^{m_i}}, c - constants \quad (10)$$

or

$$\left\| \ell_N / L_2^{(m)^*}(K_n) \right\| \leq c \cdot O(h_1^{m_1}) \cdot O(h_2^{m_2}) \cdot \dots \cdot O(h_n^{m_n}), \quad (11)$$

where

$$\ell_{N_i}(x_i) = p(x_i) \varepsilon_{[0,1]}(x_i) - \sum_{\lambda_i=1}^{N_i} C_{\lambda_i} \delta(x_i - x_i^{(\lambda_i)}),$$

$$p(x) = \prod_{i=1}^n p_i(x_i), c = \prod_{i=1}^n c_i \text{ and } m = m_1 + m_2 + \dots + m_n, m_i \geq 1, i = \overline{1, n}.$$

We are conducting proof by mathematical induction.

Suppose  $n = 2$ , then

$$x = (x_1, x_2), \quad |\alpha| = \alpha_1 + \alpha_2, \quad m = m_1 + m_2, \quad dx = dx_1 dx_2, \quad f(x) = f(x_1, x_2),$$



$p(x) = p_1(x_1) \cdot p_2(x_2)$  and  $\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2)$ .  
 If we assume in (4)  $n = 1$ , then

$$\|f_i / L_2^{(m_i)}(0, 1)\| = \left\{ \int_0^1 \left[ \frac{d^{m_i}}{dx_i^{m_i}} f(x_i) \right]^2 dx_i \right\}^{\frac{1}{2}}, (\overline{1, n}). \quad (12)$$

Thus, we have

$$\begin{aligned} |\langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle| &= |\langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \rangle| \leq \\ &\leq \| \ell_2(x_2) / L_2^{(m_2)^*}(0, 1) \| \cdot \| \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_2^{(m_1)}(0, 1) \|. \end{aligned} \quad (13)$$

Taking into account (12), (13) we compute the following norm [13]:

$$\begin{aligned} \| \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_2^{(m_1)}(0, 1) \| &= \left\{ \int_0^1 \left| \frac{d^{m_2}}{dx_2^{m_2}} \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \right|^2 dx_2 \right\}^{\frac{1}{2}} = \\ &= \left\{ \int_0^1 \left| \langle \ell_{N_1}(x_1), \frac{d^{m_2}}{dx_2^{m_2}} f(x_1, x_2) \rangle \right|^2 dx_2 \right\}^{\frac{1}{2}} \leq \\ &\leq \left\{ \int_0^1 \left[ \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \left\| \frac{d^{m_2}}{dx_2^{m_2}} f(x_1, x_2) / L_2^{(m_1)}(0, 1) \right\| \right]^2 dx_2 \right\}^{\frac{1}{2}} = \\ &= \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \left\{ \int_0^1 \int_0^1 \left[ \frac{\partial^{m_1+m_2}}{\partial x_1^{m_1} \partial x_2^{m_2}} f(x_1, x_2) \right]^2 dx_1 \right\}^{\frac{1}{2}} = \\ &= \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \left\{ \int_0^1 \int_0^1 \left[ \frac{\partial^{m_1+m_2}}{\partial x_1^{m_1} \partial x_2^{m_2}} f(x) \right]^2 dx \right\}^{\frac{1}{2}} = \\ &= c' \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \| f(x) / L_2^{(m)}(K_2) \|, c' - constants. \end{aligned} \quad (14)$$

Thus, from (13) and (14) we obtain

$$\begin{aligned} |\langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle| &\leq \\ &\leq c' \left\| \ell_{N_2}(x_2) / L_2^{(m_2)^*}(0, 1) \right\| \cdot \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \| f(x) / L_2^{(m)}(K_2) \|. \end{aligned} \quad (15)$$

Taking into account (4) from (15) we obtain

$$\| \ell_N / L_2^{(m)^*}(K_2) \| \leq c' \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \left\| \ell_{N_2}(x_2) / L_2^{(m_2)^*}(0, 1) \right\|. \quad (16)$$

Using (8), from (16) we have

$$\left\| \ell_N / L_2^{(m)^*} (K_2) \right\| \leq c' \cdot c_1 \cdot c_2 \frac{1}{N_1^{m_1} N_2^{m_2}},$$

that is

$$\left\| \ell_N / L_2^{(m)^*} (K_2) \right\| \leq c'_3 O(h_1^{m_1}) O(h_2^{m_2}), \quad (17)$$

where  $c'_3 = c' \cdot c_1 \cdot c_2$ .

Now suppose that (10) is valid for  $n = k$ , then from the above calculations we obtain [23]

$$\begin{aligned} & | \langle \ell_N(x), f(x) \rangle | = | \langle \ell_N(x_1, x_2, \dots, x_k), f(x_1, x_2, \dots, x_k) \rangle | = \\ & = | \langle \ell_{N_k}(x_k), \langle \ell_{N_{k-1}}(x_{k-1}), \dots, \langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle \dots \rangle | \leq \\ & \leq \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \left\| \ell_{N_{k-1}}(x_{k-1}) / L_2^{(m_{k-1})^*}(0, 1) \right\| \dots \\ & \cdot \left\| \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle / L_2^{(m_1)^*}(0, 1) \right\| \leq \\ & \leq \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \dots \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(x_1) \right\| \cdot c'' \left\| f(x) / L_2^{(m)}(K_k) \right\|. \end{aligned} \quad (18)$$

From (18), considering (13), we have

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq c'' \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \dots \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\|. \quad (19)$$

Then, referring to (8), from (19) we obtain

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq c'' \cdot c_1 \cdot c_2 \cdot \dots \cdot c_k \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \cdot \dots \cdot N_k^{m_k}}, \quad (20)$$

or, considering (9), from (20) we have

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq c'' \cdot d_k \cdot O(h_1^{m_1}) \dots O(h_k^{m_k}), \text{ where } d_k = \prod_{i=1}^k c_i.$$

Using the validity of assertion  $n = k$ , we prove that the assertion executed when  $n = k + 1$ . Thus, when  $n = k + 1$ , and taking account of (4) and (19), we obtain

$$\begin{aligned} & | \langle \ell_{N_{k+1}}(x_1, x_2, \dots, x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle | = \\ & = | \langle \ell_{N_1}(x_1), \langle \ell_{N_2}(x_2), \dots, \langle \ell_{N_k}(x_k), \langle \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle \dots \rangle | \leq \\ & \leq \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \cdot \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \\ & \cdot \left\| \langle \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle / L_2^{(m_{k+1})^*}(0, 1) \right\|. \end{aligned} \quad (21)$$

Using (4) and (19) from (21) we obtain

$$\begin{aligned} \left\| \ell_N / L_2^{(m)^*} (K_{k+1}) \right\| & \leq c'' \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \dots \\ & \cdot \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \left\| \ell_{N_{k+1}}(x_{k+1}) / L_2^{(m_{k+1})^*}(0, 1) \right\|. \end{aligned} \quad (22)$$

By using (8), from (22) we have

$$\left\| \ell_N / L_2^{(m)^*} (K_{k+1}) \right\| \leq c'' \cdot d_{k+1} \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \cdot \dots \cdot N_{k+1}^{m_{k+1}}},$$

where

$$d_{k+1} = \prod_{i=1}^{k+1} c_i, \quad (23)$$

or, taking into account (17), from (23) we obtain

$$\left\| \ell_N / L_2^{(m)*} (K_{k+1}) \right\| \leq c^m \cdot d_{k+1} \cdot O(h_1^{m_1}) \dots O(h_{k+1}^{m_{k+1}}), c^m \text{ constants,} \quad (24)$$

in conclusion consider, that

$$\left\| \ell_N / L_2^{(m)*} (K_n) \right\| \leq c \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_n^{m_n}}, \quad (25)$$

where  $c = c^m \cdot d_{k+1}$ .

or, considering (9), from (25), we have

$$\left\| \ell_N / L_2^{(m)*} (K_n) \right\| \leq c \cdot O(h_1^{m_1}) \dots O(h_n^{m_n}).$$

With the help of this lemma it is easy to prove the following theorem.

**Theorem 1.** The weight cubature formula (1) with the error functional (2) for  $N_1 = N_2 = \dots = N_n$ ,  $\prod_{i=1}^n N_i = N$  and  $m_1 + m_2 + \dots + m_n = m$  is optimal in order of convergence in the space  $L_2^{(m)}(K_n)$ , those, for the norm of the error functional (2) of cubature formula (1) the following holds

$$\left\| \ell_N / L_2^{(m)*} (K_n) \right\| = O(N^{-\frac{m}{n}}).$$

**Proof.** On the basis of Lemma1 under the  $N_1 = N_2 = \dots = N_n$  have  $N_i = \sqrt[n]{N}$ . Thus,

$$\prod_{i=1}^n N_i^{m_i} = N_1^{m_1 + m_2 + \dots + m_n} = N^{\frac{m}{n}}. \quad (26)$$

By substituting (26) into (25) we obtain

$$\left\| \ell_N / L_2^{(m)*} (K_n) \right\| \leq c \cdot N^{-\frac{m}{n}}. \quad (27)$$

From theorem of N.S. Bahvalov [24] and from the inequality (27) follows the proof.

## A NORM ESTIMATION FOR THE ERROR FUNCTIONAL OF WEIGHT CUBATURE FORMULAS ON THE SPACE $L_p^{(m)}(K_n)$ .

Consider the cubature formula of the form

$$\int_{K_n} p(x) f(x) dx \approx \sum_{\lambda=1}^N C_\lambda f(x^{(\lambda)}), \quad (28)$$

(28) in the Sobolev space  $L_p^{(m)}(K_n)$ , where  $K_n$  is  $n$ -dimensional unit cube. The generalized function

$$\ell_N(x) = p(x) \varepsilon_{K_n}(x) - \sum_{\lambda=1}^N C_\lambda \delta(x - x^{(\lambda)}). \quad (29)$$

is called error functional of the cubature formula (28),

$$\langle \ell_N, f \rangle = \int_{K_n} p(x) f(x) dx - \sum_{\lambda=1}^N C_\lambda f(x^{(\lambda)})$$

is an error of the cubature formula (28),  $p(x) \in L_p(K_n)$  is a weight function,  $\epsilon_{K_n}(x)$  is the characteristic function of  $K_n$ ,  $C_\lambda$  and  $x^{(\lambda)}$  are coefficients and nodes of the cubature formula (28) and  $\delta(x)$  is the Dirac delta function.

**Definition 4.** The space  $L_p^{(m)}(K_n)$  is defined as a space of functions, given on a  $n$ -dimensional unit cube  $K_n$  and having all the generalized derivatives of order  $m$ , summable with a degree  $p$  in norm [1]:

$$\|f/L_p^{(m)}(K_n)\| = \left\{ \int_{K_n} \left\{ \sum_{|\alpha|=m} \frac{m!}{\alpha!} [D^\alpha f]^2 \right\}^{\frac{p}{2}} dx \right\}^{\frac{1}{p}}, \quad (30)$$

where  $D^\alpha = \frac{\partial^m}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_n^{\alpha_n}}$ ,  $|\alpha| = \sum_{j=1}^n \alpha_j$ ,  $\alpha! = \alpha_1! \cdot \alpha_2! \cdot \dots \cdot \alpha_n!$ .

The following is true

**Lemma 2.** If for the error functional (29) of the cubature formula (28), the following conditions are fulfilled

$$\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2) \cdot \dots \cdot \ell_{N_n}(x_n),$$

where  $\ell_{N_i}(x_i) = p_i(x_i) \epsilon_{[0,1]}(x_i) - \sum_{\lambda=1}^{N_i} C_\lambda \delta(x_i - x_i^{(\lambda)})$ ,  $p(x) = \prod_{i=1}^n p_i(x_i)$  and

$$\|\ell_{N_i}/L_p^{(m_i)^*}(0,1)\| \leq d_i \frac{1}{N_i^{m_i}}, \quad d_i - \text{constants}, \quad (31)$$

that is

$$\|\ell_{N_i}/L_p^{(m_i)^*}(0,1)\| \leq d_i O(h_i^{m_i}), \quad d_i - \text{constants}, \quad (i = \overline{1, n}), \quad h_i = \frac{1}{N_i} \quad (32)$$

then

$$\|\ell_N/L_p^{(m)^*}(K_n)\| \leq d \cdot \frac{1}{\prod_{i=1}^n N_i^{m_i}}, \quad d - \text{constants}, \quad (33)$$

or

$$\|\ell_N/L_p^{(m)^*}(K_n)\| \leq d \cdot O(h_1^{m_1}) \cdot O(h_2^{m_2}) \cdot \dots \cdot O(h_n^{m_n}),$$

$d = \prod_{i=1}^n d_i$  and  $m = m_1 + m_2 + \dots + m_n$ .

**Proof.** We conduct the method of the mathematical induction. Suppose  $n = 2$   $x = (x_1, x_2)$ ,  $|\alpha| = \alpha_1 + \alpha_2$ ,  $m = m_1 + m_2$ ,  $dx = dx_1 dx_2$ ,  $f(x) = f(x_1, x_2)$ ,  $p(x) = p_1(x_1) \cdot p_2(x_2)$  and  $\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2)$ .

If presume in (30)  $n = 1$ , then

$$\|f_i/L_p^{(m_i)}(0,1)\| = \left\{ \int_0^1 \left[ \left( \frac{d^{m_i}}{dx_i^{m_i}} f(x_i) \right)^2 \right]^{\frac{p}{2}} dx_i \right\}^{\frac{1}{p}}, \quad i = 1, 2, \dots, n. \quad (34)$$

So we have

$$\begin{aligned} |\langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle| &= |\langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \rangle| \leq \\ &\leq \|\ell_{N_2}(x_2)/L_p^{(m_2)^*}(0,1)\| \cdot \|\langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_p^{(m_1)}(0,1)\|. \end{aligned} \quad (35)$$

We compute the following norm:

$$\begin{aligned}
\| \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_p^{(m_2)}(0, 1) \| &= \left\{ \int_0^1 \left[ \int_0^1 \frac{d^{m_2}}{dx_2^{m_2}} \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \right]^2 dx_2 \right\}^{\frac{1}{2}} = \\
&= \left\{ \int_0^1 \left[ \left| \langle \ell_{N_1}(x_1), \frac{d^{m_2}}{dx_2^{m_2}} f(x_1, x_2) \rangle \right|^2 \right] dx_2 \right\}^{\frac{1}{2}} \leq \\
&\leq \left\{ \int_0^1 \left[ \left( \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \cdot \left\| \frac{d^{m_2}}{dx_2^{m_2}} f(x_1, x_2) / L_p^{(m_1)}(0, 1) \right\| \right)^2 \right] dx_2 \right\}^{\frac{1}{2}} = \\
&= \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \cdot \left\{ \int_0^1 \left[ \int_0^1 \left( \frac{d^{m_1+m_2}}{dx_1^{m_1} dx_2^{m_2}} f(x_1, x_2) \right)^2 dx_1 \right] dx_2 \right\}^{\frac{1}{2}} = \\
&= d' \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \cdot \left\| f(x) / L_p^{(m)}(K_2) \right\|, \tag{36}
\end{aligned}$$

where  $d'$  - constants.

Thus, from (35) and (36) we obtain

$$\begin{aligned}
| \langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle | &\leq \\
&\leq d'' \left\| \ell_{N_2}(x_2) / L_p^{(m_2)^*}(0, 1) \right\| \cdot \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \cdot \left\| f(x) / L_p^{(m)}(K_2) \right\|. \tag{37}
\end{aligned}$$

From (37), using the definition of norm, we have

$$\left\| \ell_N(x) / L_p^{(m)^*}(K_2) \right\| \leq d'' \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \cdot \left\| \ell_{N_2}(x_2) / L_p^{(m_2)^*}(0, 1) \right\|. \tag{38}$$

Taking into account (31), on the basis of (38) we obtain

$$\left\| \ell_N / L_p^{(m)^*}(K_2) \right\| \leq d' \cdot d_1 \cdot d_2 \frac{1}{N_1^{m_1} N_2^{m_2}},$$

that is

$$\left\| \ell_N / L_p^{(m)^*}(K_2) \right\| \leq d_3 O(h_1^{m_1}) O(h_2^{m_2}), \tag{39}$$

where

$$d_3 = d' \cdot d_1 \cdot d_2.$$

Now suppose that the inequality (33) holds for  $n = k$ , then on the basis of the above calculations we obtain

$$\begin{aligned}
| \langle \ell_N(x), f(x) \rangle | &= | \langle \ell_N(x_1, x_2, \dots, x_k), f(x_1, x_2, \dots, x_k) \rangle | = \\
&= | \langle \ell_{N_k}(x_k), \langle \ell_{N_{k-1}}(x_{k-1}), \dots, \langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle \rangle \dots \rangle \rangle | \leq \\
&\leq \left\| \ell_{N_k}^{(x_k)} / L_p^{(m_k)^*}(0, 1) \right\| \cdot \left\| \ell_{N_{k-1}}(x_{k-1}) / L_p^{(m_{k-1})^*}(0, 1) \right\| \cdot \dots \\
&\dots \cdot \left\| \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle / L_p^{(m_1)}(0, 1) \right\| \leq \\
&\leq d'' \left\| \ell_{N_k}(x_k) / L_p^{(m_k)^*}(0, 1) \right\| \cdot \dots \cdot \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(x_1) \right\| \cdot \left\| f(x) / L_p^{(m)}(K_k) \right\|, \tag{40}
\end{aligned}$$

where  $d''$  - constants.

From (40), using the definition for norm of the error functional, we obtain

$$\left\| \ell_N / L_p^{(m)^*}(K_k) \right\| \leq d'' \left\| \ell_{N_1}(x_1) / L_p^{(m_1)^*}(0, 1) \right\| \dots \left\| \ell_{N_k}(x_k) / L_p^{(m_k)^*}(0, 1) \right\|. \tag{41}$$



Then, using (31), the inequality (41) reduces to

$$\left\| \ell_N / L_p^{(m)*} (K_k) \right\| \leq d'' \cdot d \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_k^{m_k}},$$

where  $d = \prod_{i=1}^k d_i$   
or

$$\left\| \ell_N / L_p^{(m)*} (K_k) \right\| \leq d'' \cdot d \cdot O(h_1^{m_1}) \dots O(h_k^{m_k}).$$

Using the validity of lemma (2) at  $n = k$ , we prove that the assertion holds for  $n = k + 1$ . Taking into account (40), at  $n = k + 1$  we estimate error of cubature formulas for the form (28)

$$\begin{aligned} & \left| \langle \ell_{N_{k+1}}(x_1, x_2, \dots, x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle \right| = \\ & = \left\| \ell_{N_1}(x_1), \langle \ell_{N_2}(x_2), \dots \langle \ell_{N_k}(x_k), \langle \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle \rangle \dots \rangle \right\| \leq \\ & \leq \left\| \ell_{N_1}(x_1) / L_p^{(m_1)*} (0, 1) \right\| \cdot \dots \cdot \left\| \ell_{N_k}(x_k) / L_p^{(m_k)*} (0, 1) \right\| \cdot \\ & \quad \cdot \left\| \langle \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle / L_p^{(m_{k+1})*} (0, 1) \right\|. \end{aligned} \quad (42)$$

Hence, as above, using the definition of norms of functionals, we get

$$\begin{aligned} \left\| \ell_N / L_p^{(m)*} (K_{k+1}) \right\| & \leq d'' \left\| \ell_{N_1}(x_1) / L_p^{(m_1)*} (0, 1) \right\| \cdot \dots \\ & \dots \cdot \left\| \ell_{N_k}(x_k) / L_p^{(m_k)*} (0, 1) \right\| \cdot \left\| \ell_{N_{k+1}}(x_{k+1}) / L_p^{(m_{k+1})*} (0, 1) \right\|. \end{aligned} \quad (43)$$

From inequalities (31) and (43) we obtain

$$\left\| \ell_N / L_p^{(m)*} (K_{k+1}) \right\| \leq d'' \cdot d_{k+1} \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_{k+1}^{m_{k+1}}}. \quad (44)$$

or

$$\left\| \ell_N / L_p^{(m)*} (K_{k+1}) \right\| \leq d'' \cdot d_{k+1} \cdot O(h_1^{m_1}) \dots O(h_{k+1}^{m_{k+1}}),$$

where  $d_{k+1} = \prod_{i=1}^{k+1} d_i$

Summarizing the results obtained, in conclude by noting that

$$\left\| \ell_N / L_p^{(m)*} (K_n) \right\| \leq d \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_n^{m_n}}. \quad (45)$$

(45) or, taking into account (32), from (44) we have

$$\left\| \ell_N / L_p^{(m)*} (K_n) \right\| \leq d \cdot O(h_1^{m_1}) \dots O(h_n^{m_n}), \quad d - \text{constant}.$$

Lemma 2 is proved.

With the help of this lemma it is easy to prove the following theorem.

**Theorem 2.** The weight cubature formula (28) with the error functional (29) at  $N_1 = N_2 = \dots = N_n$ ,  $\prod_{i=1}^n N_i = N$  and  $m_1 + m_2 + \dots + m_n = m$  is optimal in order of convergence in the space  $L_p^{(m)}(K_n)$ , for the norms of error functional (29) of the cubature formula (28) have the equality

$$\left\| \ell_N / L_p^{(m)*} (K_n) \right\| = O\left(N^{-\frac{m}{n}}\right).$$

**Proof.** On the basis of Lemma 2 at  $N_1 = N_2 = \dots = N_n$  we have  $N_i = \sqrt[n]{N}$ ,  $i = 1, 2, \dots, n$ . Thus,

$$\prod_{i=1}^n N_i^{m_i} = N_1^{m_1+m_2+\dots+m_n} = N^{\frac{m}{n}}. \quad (46)$$

By substituting (46) into inequality (45), we obtain

$$\left\| \ell_N / L_p^{(m)*}(K_n) \right\| \leq C \cdot N^{-\frac{m}{n}}. \quad (47)$$

From the N.S. Bakhvalov theorem [24] and the inequality (47) follows the proof of the theorem.

## WEIGHT CUBATURE FORMULAS IN THE SPACE $L_2^{(m)}(K_n)$

Multidimensional cubature formulas differ from the one-dimensional with two features:

- 1) infinitely varied forms of multidimensional areas of integration;
  - 2) rapidly grows number of integration nodes with increasing space dimension.
- Problem 2) requires special attention to the construction of the most efficient formulas.

Here, we discuss the formula with taking into account this requirement. It is known, that such formulas are called by N.S. Bakhvalov as "practical formulas" [24].

We regard the weight cubature formula

$$\int_{\Omega} p(x) f(x) dx \approx \sum_{\lambda=1}^N C_{\lambda} f(x^{(\lambda)}), \quad (48)$$

in space  $L_2^{(m)}(K_n)$ , where  $K_n$  -  $n$  - dimensional unit cube and  $p(x) \in L_2(K_n)$  is weight function. In the cubature formula (48) comparable to generalized function

$$\ell_N(x) = p(x) \varepsilon_{K_n}(x) - \sum_{\lambda=1}^N C_{\lambda} \delta(x - x^{(\lambda)}). \quad (49)$$

and we call it a error functional.

**Definition 5.** The space  $L_2^{(m)}(K_n)$  is defined as the space of functions, given on the  $K_n$  and the norm of a function, which is determined by the following equation

$$\|f / L_2^{(m)}(K_n)\| = \left\{ \int_{K_n} \left( \frac{\partial^m f(x)}{\partial x_1^{m_1} \partial x_2^{m_2} \dots \partial x_n^{m_n}} \right)^2 dx \right\}^{\frac{1}{2}}, \quad (50)$$

where  $m_1 + m_2 + \dots + m_n = m$ ,  $m_i \geq 0$ ,  $i = \overline{1, n}$   
with the scalar product

$$(f, \varphi)_{L_2^{(m)}(K_n)} = \int_{K_n} \left( \frac{\partial^m f(x)}{\partial x^m} \right) \left( \frac{\partial^m \varphi(x)}{\partial x^m} \right) dx,$$

where  $\partial x^m = \partial x_1^{m_1} \partial x_2^{m_2} \dots \partial x_n^{m_n}$ ,  $m = m_1 + m_2 + \dots + m_n$ ,  $dx = dx_1 dx_2 \dots dx_n$ .

As it is known [1], the norm of a function in the space  $L_2^{(m)}(K_n)$  determined by the formula

$$\|f / L_2^{(m)}(K_n)\| = \left\{ \int_{K_n} \sum_{|\alpha|=m} \frac{m!}{\alpha!} (D^{\alpha} f(x))^2 dx \right\}^{\frac{1}{2}}, \quad (51)$$

where  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ ,  $\alpha! = \alpha_1! \cdot \alpha_2! \cdot \dots \cdot \alpha_n!$  and  $D^\alpha f(x) = \frac{\partial^{|\alpha|} f(x_1, \dots, x_n)}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_n^{\alpha_n}}$ .

Suppose that in (51)  $n = 2$  and  $m = 2$ , then we obtain the following

$$\begin{aligned} \int_{K_2} \sum_{|\alpha|=m} \frac{m!}{\alpha!} \left( \frac{\partial^m f(x)}{\partial x^m} \right)^2 dx &= \int_{K_2} \sum_{\alpha_1+\alpha_2=2} \frac{2!}{\alpha_1! \alpha_2!} \left( \frac{\partial^2 f(x)}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2}} \right)^2 dx = \\ &= \int_{K_2} \left[ \left( \frac{\partial^2 f(x)}{\partial x_2^2} \right)^2 + \frac{2!}{1! \cdot 1!} \left( \frac{\partial^2 f(x)}{\partial x_1 \partial x_2} \right)^2 + \left( \frac{\partial^2 f(x)}{\partial x_1^2} \right)^2 \right] dx. \end{aligned} \quad (52)$$

When  $n = 2$  and  $m = 2$  equality (50) takes the following form:

$$\|f/L_2^{(2)}(K_2)\|^2 = \int_{K_2} \left( \frac{\partial^2 f(x)}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2}} \right)^2 dx. \quad (53)$$

Obviously, in the right hand side of (53) is less computing, than in (52), and it follows that the norm of the function in space  $L_2^{(2)}(K_2)$  the number of computing operations will be much less, than in space  $L_2^{(2)}(K_2)$ , as in the norm (53), involved only the mixed derivatives. Now we prove the the following theorem, which is one of the main results of this work.

**Theorem 3.** If, for the error functional (49) of the weight substure formula (48) in the space  $barL_2^{(m)}(K_n)$  the following conditions are fulfilled

$$\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2) \cdot \dots \cdot \ell_{N_n}(x_n)$$

and

$$\|\ell_{N_i}/L_2^{(N_i)}(0,1)\| \leq d_i \frac{1}{N_i^{m_i}}, \quad d_i - constants \quad (54)$$

that is

$$\|\ell_{N_i}/L_2^{(m_i)}(0,1)\| \leq d_i O(h_i^{m_i}), \quad d_i - constants, (i = \overline{1, n}), h_i = \frac{1}{N_i} \quad (55)$$

then

$$\|\ell_N/L_2^{(m)}(K_n)\| \leq d \cdot \frac{1}{\prod_{i=1}^n N_i^{m_i}}, \quad d - constants, \quad (56)$$

or

$$\|\ell_N/L_2^{(m)}(K_n)\| \leq d \cdot O(h_1^{m_1}) \cdot O(h_2^{m_2}) \dots \cdot O(h_n^{m_n}), \quad (57)$$

where

$$\ell_{N_i}(x_i) = p_i(x_i) \varepsilon_{[0,1]}(x_i) - \sum_{k_i=1}^{N_i} C_{k_i} \delta(x_i - x_i^{(k_i)}), \quad p(x) = \prod_{i=1}^n p_i(x_i),$$

$d = \prod_{i=1}^n d_i$ ,  $m = m_1 + m_2 + \dots + m_n$  and  $m_i$  is arbitrary ( $i = \overline{1, n}$ ), and  $m_i \geq 1$ .

**Proof.** We are conducting proof by mathematical induction.

Suppose  $n = 2$ , then

$x = (x_1, x_2)$ ,  $|\alpha| = \alpha_1 + \alpha_2$ ,  $m = m_1 + m_2$ ,  $dx = dx_1 dx_2$ ,  $f(x) = f(x_1, x_2)$ ,

$p(x) = p_1(x_1) \cdot p_2(x_2)$  and  $\ell_N(x) = \ell_{N_1}(x_1) \cdot \ell_{N_2}(x_2)$ .  
If presume in (50)  $n = 1$ , then

$$\|f_i / L_2^{(m_i)}(0, 1)\| = \left\{ \int_0^1 \left( \frac{\partial^{m_i} f(x_i)}{\partial x_i^{m_i}} \right)^2 dx_i \right\}^{\frac{1}{2}}, (i = \overline{1, n}).$$

Thus, we have

$$|\langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle| = |\langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \rangle| \leq \| \ell_{N_2}(x_2) / L_2^{(m_2)*}(0, 1) \| \cdot \| \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_2^{(m_1)*}(0, 1) \|. \quad (58)$$

We compute the following norm:

$$\begin{aligned} \| \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle / L_2^{(m_1)*}(0, 1) \| &= \left\{ \int_0^1 \left| \frac{\partial^{m_1}}{\partial x_1^{m_1}} \langle \ell_{N_1}(x_1), f(x_1, x_2) \rangle \right|^2 dx_1 \right\}^{\frac{1}{2}} = \\ &= \left\{ \int_0^1 \left| \langle \ell_{N_1}(x_1), \frac{\partial^{m_1}}{\partial x_1^{m_1}} f(x_1, x_2) \rangle \right|^2 dx_1 \right\}^{\frac{1}{2}} \leq \\ &\leq \left\{ \int_0^1 \left[ \left\| \ell_{N_1}(x_1) / L_2^{(m_1)*}(0, 1) \right\| \cdot \left\| \frac{\partial^{m_1}}{\partial x_1^{m_1}} f(x_1, x_2) / L_2^{(m_1)}(0, 1) \right\| \right]^2 dx_1 \right\}^{\frac{1}{2}} = \\ &= \left\| \ell_{N_1}(x_1) / L_2^{(m_1)*}(0, 1) \right\| \cdot \left\{ \int_0^1 \left[ \int_0^1 \left[ \frac{\partial^{m_1+m_2}}{\partial x_1^{m_1} \partial x_2^{m_2}} f(x_1, x_2) \right]^2 dx_1 \right] dx_2 \right\}^{\frac{1}{2}} = \\ &= \left\| \ell_{N_1}(x_1) / L_2^{(m_1)*}(0, 1) \right\| \cdot \| f(x) / L_2^{(m)}(K_2) \|. \quad (59) \end{aligned}$$

where  $x = (x_1, x_2)$  and  $m = m_1 + m_2$ .  
Thus, from (58) and (59) we obtain

$$\begin{aligned} |\langle \ell_N(x_1, x_2), f(x_1, x_2) \rangle| &\leq \| \ell_{N_2}(x_2) / L_2^{(m_2)*}(0, 1) \| \cdot \\ &\cdot \| \ell_{N_1}(x_1) / L_2^{(m_1)*}(0, 1) \| \cdot \| f(x) / L_2^{(m)}(K_2) \|. \quad (60) \end{aligned}$$

Considering (50) from (60) we obtain

$$\| \ell_N / L_2^{(m)*}(K_2) \| \leq \| \ell_{N_1}(x_1) / L_2^{(m_1)*}(0, 1) \| \cdot \| \ell_{N_2}(x_2) / L_2^{(m_2)*}(0, 1) \|. \quad (61)$$

Taking into account (54) from (61) we have

$$\| \ell_N / L_2^{(m)*}(K_2) \| \leq d_1 \cdot d_2 \cdot \frac{1}{N_1^{m_1} \cdot N_2^{m_2}},$$

or

$$\left\| \ell_N / L_2^{(m)^*} (K_2) \right\| \leq d_3' O(h_1^{m_1}) \cdot O(h_2^{m_2}), \quad (62)$$

where  $d_3' = d_1 \cdot d_2$ .

When  $n = k$  we obtain

$$\begin{aligned} & | \langle \ell_N(x), f(x) \rangle | = | \langle \ell_N(x_1, x_2, \dots, x_k), f(x_1, x_2, \dots, x_k) \rangle | = \\ & | \langle \ell_{N_k}(x_k), \langle \ell_{N_{k-1}}(x_{k-1}), \dots \langle \ell_{N_2}(x_2), \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle \rangle \dots \rangle \rangle | \leq \\ & \leq \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \left\| \ell_{N_{k-1}}(x_{k-1}) / L_2^{(m_{k-1})^*}(0, 1) \right\| \dots \\ & \cdot | \langle \ell_{N_1}(x_1), f(x_1, x_2, \dots, x_k) \rangle / L_2^{(m_1)}(0, 1) | \leq \\ & \leq \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \dots \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(x_1) \right\| \cdot \left\| f(x) / L_2^{(m)}(K_k) \right\|. \end{aligned} \quad (63)$$

From (63), taking into account (50) we have

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \dots \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\|. \quad (64)$$

Then in view of (54) from (64) we obtain

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq d \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \cdot \dots \cdot N_k^{m_k}}, \quad \text{where } d = \prod_{i=1}^k d_i, \quad (65)$$

or, taking into account (55), from (65) we will have

$$\left\| \ell_N / L_2^{(m)^*} (K_k) \right\| \leq d \cdot O(h_1^{m_1}) \dots O(h_k^{m_k}).$$

Using validity of the assertion of theorem 1 for  $n = k$ , we prove that the assertion is performed when  $n = k + 1$ .

Thus, suppose  $n = k + 1$ , then taking into account (50), from (64) we have

$$\begin{aligned} & | \langle \ell_{k+1}(x_1, x_2, \dots, x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle | = \\ & = | \langle \ell_{N_1}(x_1), \langle \ell_{N_2}(x_2), \dots \langle \ell_{N_k}(x_k), \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle \dots \rangle \rangle | \leq \\ & \leq \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \dots \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \\ & \cdot \left| \langle \ell_{N_{k+1}}(x_{k+1}), f(x_1, x_2, \dots, x_{k+1}) \rangle / L_2^{(m_{k+1})}(0, 1) \right| \end{aligned} \quad (66)$$

Using (50) and (64) from (66) we obtain

$$\begin{aligned} & \left\| \ell_N / L_2^{(m)^*} (K_{k+1}) \right\| \leq \left\| \ell_{N_1}(x_1) / L_2^{(m_1)^*}(0, 1) \right\| \dots \\ & \cdot \left\| \ell_{N_k}(x_k) / L_2^{(m_k)^*}(0, 1) \right\| \cdot \left\| \ell_{N_{k+1}}(x_{k+1}) / L_2^{(m_{k+1})^*}(0, 1) \right\|. \end{aligned} \quad (67)$$



By using (54) from (67) we have

$$\left\| \ell_N / L_2^{(m)*}(K_{k+1}) \right\| \leq d_{k+1} \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_{k+1}^{m_{k+1}}}, \quad (68)$$

or, considering (62) and (68) we obtain

$$\left\| \ell_N / L_2^{(m)*}(K_{k+1}) \right\| \leq d_{k+1} \cdot O(h_1^{m_1}) \dots O(h_{k+1}^{m_{k+1}}), \text{ where } d_{k+1} = \prod_{i=1}^{k+1} d_i.$$

Thus obtained inequalities (56) and (57):

$$\left\| \ell_N / L_2^{(m)*}(K_n) \right\| \leq d \frac{1}{N_1^{m_1} \cdot N_2^{m_2} \dots N_n^{m_n}}, \quad d - \text{constants}, \quad (69)$$

or, considering (55) from (69) we obtain

$$\left\| \ell_N / L_2^{(m)*}(K_n) \right\| \leq d \cdot O(h_1^{m_1}) \dots O(h_n^{m_n}), \quad h_l = \frac{1}{N_l}, \quad l = \overline{1, n}, \quad (70)$$

where  $d = \prod_{i=1}^n d_i$ .

If, in (69) or (70) suppose  $N = N_1 \cdot N_2 \cdot \dots \cdot N_n$ ,  $N_1 = N_2 = \dots = N_n$  and  $m_1 + m_2 + \dots + m_n = m$  then we have

$$\left\| \ell_N / L_2^{(m)*}(K_n) \right\| \leq d \cdot N^{-\frac{m}{n}}$$

or

$$\left\| \ell_N / L_2^{(m)*}(K_n) \right\| \leq d \cdot O(h^m), \quad d - \text{constant}, \quad h = N^{-\frac{1}{n}}. \quad (71)$$

which was needed of proof. Theorem is proved.

Thus, we obtain an upper estimate for the norm of the error functional (49) for cubature formula (48) in the space  $L_2^{(m)*}(K_n)$ .

A similar assessment was obtained previously for the norm of the error functional of cubature formula (48) on the quotient space of S.L. Sobolev  $L_2^{(m)}(K_n)$  and as a result we have received the same order of convergence to zero as  $N \rightarrow \infty$ , although the norm of function was defined in different ways, this is confirmed by the inequality (27), (71).

For illustration, we present an example at  $n = 2$ . Suppose

$$(x_1, x_2) = e^{x_1} \left( \frac{1}{2} - x_2^2 \right)^{3/2}. \quad (72)$$

Obviously, that the derivatives

$\frac{\partial^{m-1} f(x_1, x_2)}{\partial x_1^{m-1}}$  and  $\frac{\partial f(x_1, x_2)}{\partial x_2}$  are continuous on  $K_2$ , but  $\frac{\partial^2 f(x_1, x_2)}{\partial x_2^2}$  has a feature on  $K_2$ . Therefore, from the condition  $m = m_1 + m_2$  it is clear that  $m_1 = m - 1$  and  $m_2 = 1$ , in that  $m - 1 + 1 = m$ .

Hence it follows that  $f(x_1, x_2) \in L_2^{(m)}(K_2)$  when  $m_1 = m - 1$ ,  $m_2 = 1$  and  $f(x_1, x_2) \notin L_2^{(m)}(K_2)$ .

## CONCLUSION

In this paper we investigate weight cubature formula in function spaces of S.L. Sobolev  $L_2^{(m)}$ ,  $L_p^{(m)}$ ,  $L_2^{(m)}$  for the functions defined in the  $n$  - dimensional unit cube  $K_n$  and obtain an upper estimate for the norm of error functionals of weight cubature formulas. The basis of theorem N.S. Bahvalov it is proved that considered viewed cubature formulas are optimal on order of convergence in these spaces.

## REFERENCES

1. S. L. Sobolev, *Introduction to the theory of cubature formulas* (Moscow, Nauka, 1974).
2. M. D. Ramazanov, *Lectures on theory of approximate integration* (Bashkir State University, 1973).
3. G. N. Salikhov, *Cubature formulas for multidimensional sphere*. (Tashkent, Fan, 1985).
4. K. M. Shadimetov, "Construction of weight optimal quadrature formulas in  $L_2^{(m)}(0, 1)$ ," *Siberian journal of Computational Mathematics*, Novosibirsk. **5**, 275–293 (2002).
5. K. M. Shadimetov, *Layitice quadrature and cubature formulas in the Sobolev space. Dissertation of Doctor of Sciences*, PhD dissertation, Tashkent (2002), a full PHDTHESIS entry.
6. T. Catinas and G. Coman, "Stability of nonlinear modes," *Mathematica* **52**, 1–16 (2005).
7. P. Blaga and G. Coman, "Some problems on optimal quadrature. stadia univ," *Mathematica* **52**, 21–44 (2007).
8. P. Kohler, "On the weights of sard's quadrature formulas," *Calcolo* **25**, 169–186 (1988).
9. F. Lanzara, "On optimal quadrature formulae," of *Inequality Appl* **5**, 201–225 (2000).
10. Kh.M.Shadimetov and A.R.Hayotov, "Optimal quadrature formulas with positive coefficients in  $L_2^{(m)}(0, 1)$ ," *Journal of Computational and Applied Mathematics* **235**, 1114–1128 (2011).
11. A. R. Hayotov, G. Milovanovic, and K. M. Shadimetov, "On an optimal quadrature formula in the sense of sard," *Numerical Algorithms* **57**, 487–510 (2011).
12. T. H. Sharipov, *Some problems of the theory of approximate integration. Dissertation of Candidate of Sciences*, PhD dissertation, Tashkent (1975), a full PHDTHESIS entry.
13. Kh.Shadimetov and F. A. Nuraliev, "Optimization of quadrature formulas with derivatives," *Problems of computational and applied mathematics* **4**, 61–70 (2017).
14. A.Nuraliev, "Minimisation the error functionals norm of the hermitian type quadrature formula," *Uzbek Mathematical journal* **1**, 53–64 (2015).
15. Kh.M.Shadimetov and B.S.Daliev, "The extremal function of quadrature formulas for the approximate solution of the generalized abel integral equation," *Problems of computational and applied mathematics* **2**, 88–96 (2019).
16. A. R. Hayotov and S. S. Babaev, "Optimal quadrature formulas for computing of Poisson integrals in a Hilbert space," *Problems of computational and applied mathematics* **4(28)**, 73–84 (2020).
17. O.I.Jabolov, "Higher bound for the norm of the error functional of weight cubature formulas in the space," *Modern problems of the applied mathematics and information technology - Al- Khorezmiy* **2012** **1**, 19–22 (2012).
18. O.I.Jabolov, "The bound of the weight cubature formulas in the space," *Applied mathematics and information security*, 25–30 (2014).
19. O.I.Jabolov, "Counting the norm of the error functional of the interpolation formulas periodical function in the space sobolev," *Problems of computational and applied mathematics* **4**, 53–58 (2015).
20. O.I.Jabolov, "The optimal in the order of weight cubature formulas of the hermitian type in the space sobolev," *East European Scientific Journal* **85/21**, 162–170 (2016).
21. O.I.Jabolov, "Higher bound the norm of the error functional of weight cubature formulas of the hermitian type in the space sobolev," *Problems of computational and applied mathematics* **3**, 70–78 (2017).
22. O.I.Jabolov, "The lower bound for the norm of the error functional of lattice cubature formulas in the space," *Modern problems of the applied mathematics and information technology - Al- Khorezmiy* **2018** **1**, 149–150 (2018).
23. S. L. Sobolev, *Some applications of functional analysis in mathematical physics* (Leningrad, Nauka, 1950).
24. N. S. Bahvalov, *Numerical methods* (Moscow, Nauka, 1973).
25. A. R. Hayotov, G. V. Milovanović, and Kh. M. Shadimetov, "Optimal quadratures in the sense of Sard in a Hilbert space," *Applied Mathematics and Computation* **259**, 637–653 (2015).
26. Kh. M. Shadimetov, A. Hayotov, and F. A. Nuraliev, "On an optimal quadrature formula in sobolev space  $L_2^{(m)}(0, 1)$ ," *J. Comput. Appl. Math.* **243**, 91–112 (2013).
27. S. S. Babaev and A. R. Hayotov, "Optimal interpolation formulas in the space  $W_2^{(m,m-1)}$ ," *Calcolo* **56**, doi.org/10.1007/s10092-019-0320-9 (2019).
28. O. S. Zikirov, "Boundary-value problems for a class of third-order composite type equations," *Operator Theory: Advances and Applications* **216**, 331–341 (2011).

