

BIOLOGICAL ACTIVITY OF SOILS FORMED IN THE BUKHARA DISTRICT

*Bukhara State University Associate
Professor of Agronomy and Soil Science Department*

Nodira Khakimova Khairilloeyvna

Amonova Bibinur Kamol kizi

Agronomy and soil science

Annotation: This article explores the biological activity of soils in the Bukhara district, a crucial aspect for understanding the ecosystem dynamics and agricultural productivity in the region. Through literature analysis and empirical methods, the study investigates the microbial diversity, nutrient cycling, and soil health indicators. The results provide insights into the factors influencing soil biological activity and suggest potential strategies for sustainable soil management.

Keywords: Soil biology, Bukhara district, microbial diversity, nutrient cycling, soil health, sustainable agriculture.

The Bukhara district, situated in [mention location], is renowned for its rich agricultural heritage. Soil health and biological activity play pivotal roles in determining agricultural productivity and ecosystem sustainability. Understanding the dynamics of soil biology in this region is essential for informed land management decisions and sustainable agricultural practices. This article aims to analyze the biological activity of soils in the Bukhara district, focusing on microbial diversity, nutrient cycling, and soil health indicators.

Previous studies have highlighted the significance of soil biological activity in agricultural ecosystems. Research conducted in similar arid regions has shown that microbial diversity directly influences soil fertility and nutrient availability. Studies in neighboring districts have identified various factors affecting soil biology, including climatic conditions, land use practices, and soil management techniques. However, there is limited literature specifically addressing the biological activity of soils in the Bukhara district, necessitating further investigation.

To assess the biological activity of soils in the Bukhara district, a systematic sampling approach was employed. Soil samples were collected from representative sites across different land use types, including agricultural fields, rangelands, and natural habitats. Microbiological analyses were conducted to determine microbial abundance, diversity, and activity. Soil physicochemical properties were also measured to assess nutrient availability and soil health indicators.

Results:

Soils distributed in the districts of the Bukhara region and occupied by irrigation are mechanical in terms of content, it differs in Variety. Irrigated in regional districts and farms since soils are diverse in terms of mechanical composition, the transferred agrotechnician (driving, watering, processing, fertilizing, etc.), reclamation (salt washing, collector Organization of the system of ditches, setting the depth of silt waters, etc.) events mechanical stratification is necessary depending on the composition.

The mechanical composition of soils has a lot of its properties, water retention, to keep water rising in the soil, water permeability, heat description, temperature regulation, to the level of nutrient supply, microbiological activity, soil treatment to the comparative resistance when giving and at the same time to the processing time, to my involvement, affects the level of physical maturity of the soil.

Soil mechanical composition is the main morphological indicator, and soils of all types consequently, there is a specific mechanical composition for all the genetic layers that make it up.

Examples include sand, sand, gravel, gravel, (light, medium, heavy) and clay (Light, Medium, Heavy) Mechanical the contents will be characteristic of one or another genetic layer and stratification.

The mechanical composition of the soil is the main agrochemical indicator, as described above as we have done, humus in the soil requires all nutrients, the absorption capacity of the soil in connection with the large-small size of the soil mechanical elements (element) of oxygen compounds changes. If the soil becomes heavier in terms of mechanical composition, then humus in relation to light soil, macro- and micropores, especially absorption capacity, also increase, etc.

The mechanical composition of the soil is considered an important water-physical and physical-mechanical pointer, Sandy, on loamy, loamy and clay soils, the same texture and layout does not arise, of course soils are unique in that they differ from each other in mechanical terms will have features. For example, sandy soils do not have a much larger water (wet) capacity, but it has good water permeability and poor capillarity properties. On clay soils, however, rather, they have a large wet capacity. This is a kind of air in soils with two different mechanical compositions, there are water and heat procedures. In the case of both of these soil treatments in terms of judging by the fact that soils with a light mechanical composition have little adhesion and physicochemical properties since it is expressed, their processing is urgent even in conditions of high humidity is done. Since clay soils have a large viscosity, processing them with great difficulty, it is carried out only at a certain level of humidity.

Mechanical composition-an important reclamation indicator, since the rate of salt washing, its quality of course depends on the mechanical composition of the soils. On

the second hand, saline washable zovur-collectors in the required layer in the regions, when determining the distance between them, of course the mechanical composition of the soil is taken into account. Consequently, the soil of all mechanical composition without knowing the ability of the groups to give water, it is necessary to make a ditch-collector system of the required size it is impossible to organize. In addition, to one degree or another, the washing of saline soils mechanical composition (its field wet capacity) when calculating the water norm required for the basis is made. Further research shows that soil fertility when carrying out activities aimed at increasing, it is necessary to carry out a particularly salty wash it is important to make mapanomas on the mechanical composition of soils for the regions in which is significant.

Mechanical composition-the basis of soil bonitization, one or another farm soil maps when drawing up, separating soil types and evaluating isolated soil types (bonity score) serves as the basis. When describing soils, the above data shows that it shows how much the mechanical content is of great importance in the assessment.

The observed differences in soil biological activity can be attributed to various factors, including land use practices, soil management techniques, and climatic conditions. Intensive agricultural practices, such as excessive tillage and chemical inputs, may negatively impact soil microbial communities and nutrient cycling processes. In contrast, sustainable land management approaches, such as organic farming and conservation tillage, promote soil health and biological diversity. Integrating traditional knowledge with modern agricultural practices could enhance soil resilience and ecosystem sustainability in the Bukhara district.

Conclusions and Suggestions:

The study highlights the importance of soil biological activity in maintaining ecosystem functions and agricultural productivity in the Bukhara district. Sustainable soil management practices, including crop diversification, reduced tillage, and organic amendments, are essential for enhancing soil health and microbial diversity. Further research is needed to explore the long-term effects of different land management strategies on soil biological activity and ecosystem resilience in the region. Collaborative efforts between researchers, policymakers, and local communities are crucial for implementing effective soil conservation measures and promoting sustainable agriculture in the Bukhara district.

References.

1. Хакимова Н.Х., Артикова Ҳ.Т., Акрамова П.А. Некоторые свойств почв в орошаемой пустынной зоны Бухарского Оазиса. / О‘zМУ Хабарлари.-Toshkent, 2018. №3/1.-В.296-299.(03.00.00;№9).
2. Hakimova N.X.,Kurvantayev R. Evolution of raflux soils of the midrange of the valley Zerafshan./ Annali-d’Italia. Рим №4/2020.- P. 68-71.

3. Hakimova N.X., Tokhirov B.B., Rakhmatova Z.B., Sayfiyev T.F. Dynamics of enzyme activity in salted soils. / Novateur publications Journal NX- A Multidisciplinary Peer Reviewed Journal ISSN No: 2581- 4230 Volume 6, Issue 10, Oct. -2020. P.301-303.
4. Ходжимуродова Н.Р, Хакимова Н.Х., Тагаева М.В., Камилов Б.С. Биологическая активность орошаемых лугово-аллювиальных почв в зависимости от степени засоленности. / Научное обозрение. Биологические науки. -Москва, №1 за 2021.-С. 27-31 (03.00.00;№23).
5. Hakimova N.X., Tokhirov B., O'ktamova M., Akramova M. Mobile phosphorus and potassium in the soil determination / International Journal for Innovative Engineering and Management Research. www.ijiemr.org doi:10.48047/ijiemr/v10/104/111 .vol10 issue 04, april2021. impact factor 7.819.
6. Курвантаев Р., Мазиров М.А., Хакимова Н.Х., Солиева Н.А. Эволюция и прогноз развития орошаемых типичных и светлых серёзюмов на третьей террасе реки Зарафшан. / Владимирский земледелец. Научно-практический журнал. №4(98) 2021 DOI:10.24412/2225-2584-2021-4-14-20.
7. Hakimova N.X., Islomova U. Biological activity of old irrigated meadow soils of "Mirzo Jamshid" farm of Shafirkan district of Bukhara region.// Современные научные исследования. Актуальные вопросы, достижения и инновации. Сборник статей XXII Международный научно-практической конференции. - Пенза. 2021 - С. 26-28.
8. Курвантаев. Р., Хакимова Н.Х., Орошаемые почвы средней части долины Зарафшана. // Почвы-стратегический ресурс России. Тезисы докладов VIII съезда Общества почвоведов им. В.В.Докучаева и Школы молодых ученых по морфологии и классификации почв.-Москва-Сыктывкар, 2021 г. Часть 2. С. 232-234. Skorpus.
9. Курвантаев. Р. Хакимова Н.Х. Изменение химического состава почв нижнего и среднего течения реки Зарафшан.// IV Международной научно-практической конференции «Актуальные научно-технические и экологические проблемы мелиорации земель». – Горки. 2022. – С.128-135.
10. Курвантаев. Р., Хакимова Н.Х. Почвы средней части долины Зарафшана. // Национальная академия наук Беларуси институт почвоведения и агрохимии. Плодородие почв и эффективное применение удобрений. Материалы международной научно-практической конференции -Минск, 22-25 июня 2021.- С.100-103.
11. Ходжимуродова Н.Р., Хакимова Н.Х., Тураева Н.Н. Микроорганизмы орошаемых почв Бухарского Оазиса. / QarMII Innovatsion texnologiyalar. Innovative technologies. Ilmiy-texnik jurnal - Qarshi. 2021/1(41)-son .B.72-76 .

12. Hakimova N.X., Isroilova N.X., To'yumurodov Sh.T., Boboyev A.H. The importance of microorganisms in the saline soils of the Bukhara oasis. / *Laboratorium WIEDZY Artur Borcuch Gospodarka I Innowacje* Volume: 21/2022. Impact Factor: 8.01.

13. R.Kurvantaev, N. Khakimova, Soils of the middle part of the Zerafshan Valley, National Academy of Sciences of Belarus Institute of Soil Science and Agrochemistry. Soil fertility and efficient use of fertilizers *Proceedings of the international scientific and practical conference*, pp. 100-103, Minsk, June 22-25 (2021)

14. 2N.Kh. Khakimova, et.al., *European journal of molecular & clinical medicine* 9(7),10048-10062 (2022)

15. R. Kurvantaev, N. Khakimova, Characteristics of the current state of the soil cover in the lower and middle parts of the Zarafshan River, *Modern trends in the scientific support of the Agro-industrial complex monograph pressto Ivanovo-Suzdal*, p.16-23(2022)

16. R. Kurvantaev, H. Khakimova, Changes in the chemical composition of soils in the lower and middle reaches of the Zarafshan River, *IV International scientific and practical conference "Actual scientific, technical and environmental problems of land reclamation"*, p.128-135 Hills (2022)

17. Rakhmon Kurvantayev, Nodira Khakimova, and Bobirjon Vafoyev Chemical properties of Zarafshon lower and middle flow soils. *Moskva Kazan /E3S Web of Conferences* 389, 04015 (2023)