

Use of *Pistia Stratiotes* (Araceae) in Water Quality Management in Khorezm Region (Uzbekistan)

¹Kupalboy Yuldashev, ²Ikram Abdullaev, ³Sulaymon Buriyev, ⁴Zokir Tajiyev

¹Urgench State University, Urgench, Uzbekistan.

²Khorezm Academy of Mamun, Khiva, Uzbekistan.

³Bukhara State University, Bukhara, Uzbekistan.

⁴Urgench State University, Urgench, Uzbekistan.

E-mail:kopalboyoldashev@gmail.com

E-mail:a_ikrom@mail.ru

Abstract: In this article there is described about Khorezm region conditions *Pistia* water cabbage (*Pistia stratiotes*) scientific and practical significance and methods of plant biology, reproduction in biological pools and their use in biological treatment of waste water. The results of the study are mentioned. Research work was carried out in flowing water of the effluent from the Khiva Carpet plant in Khiva, based on the cultivation of the *pistiya* plant. When studying the physicochemical composition of wastewater in the mouth, the temperature of the experimental water rose from 27.50C to 33.80C, the water (Ph) medium increased from acidic to alkaline (6.8-7.6), the color changed from reddish-yellow to colorless, the odor disappeared, decreased amount of suspended solids (107.5-72.5 mg / l), increased amount of oxygen (1.4-12.3 mg / l), decreased amount of K₂S₂O₈ (123.5-28.7 mg / l), During the oxidation process, the amount of O₂ decreased from 118.4 mg / l to 28.9 mg / l and the amount of ammonia to 9.0-1.8 mg / l, as well as the loss of nitrites, nitrates and sulfates 121.0-67.0 mg / l, chlorides 113.0-54.8 mg / l, phosphates 8.3-1.4 mg / l and it was determined that this water could be reused.

Keywords: *pistiya*, biological method, biological pool, waste water, biomass allocation, growth analysis, population ecology

Introduction

Pistiastratiotes Linnaeus (Araceae) (water lettuce) is a free floating macrophyte capable of rapid vegetative propagation (Täckholm 1974). The morphological structure of *P. stratiotes* has caused it to be one of the most notorious weeds. *Pistiastratiotes* were listed as an




Fig.1. Map area of field survey (Urgench district, tugai forest Cholish)

The area is a lowland plant located in the NW part of Uzbekistan, along the lower reaches of the Amu darya river, between 600C-610C longitude and 410C -420C latitude, at 113-138 m above sea level. The climate is extremely continental, with an average annual precipitation of 80-90 mm. Average temperature in January is -50C, in July + 300C. Meadow, meadow marshly, marsh-sandy an typical alkali soils prevailThe climate of the oasis is greatly influenced by the deserts of Kyzylkum and Karakum (Abdullaev et al., 2019; Abdullaev, et al., 2020).

2.2 Plant biomass

The plant materials from each square were separated into leaves, stem-bases, reproductive structures, stolons and roots. Afterwards, the dry weight was estimated after oven drying at 85 ° C until constant weight. All biomass measures were determined as grams of dry matter per square meter (g DM m²). Total biomass (g DM m²) was calculated and divided by the total density (individual m²) to give the average dry weight of the individual (g DM individual¹)The proportional biomass allocation was calculated as the biomass of a specific organ divided by the total biomass (Bazzaz & Grace 1997).

2.3 Water sampling and analysis

Physicochemical changes in wastewater during the study, ie in determining the composition of wastewater before and after sowing of higher aquatic plants (pistachios) Lure (1984), Strogonov (1980) methods, as well as geobotanical observation and biochemical methods. Each month, three

(Eichornia), azolla (*Azolla microphylla*) such as high aquatic plants.

The following results can be achieved in the process of biological treatment of wastewater:

- Sewage is cleaned of biogenic salts, ammonia, nitrite, nitrate, sulfates, chlorides, oil residues are reduced. As a result of photosynthesis, the amount of oxygen in the water increases several times. The wastewater is cleaned of pathogenic bacteria.
- The suspended solids in the water are retained and assimilated and purified. The amount of water-insoluble substances is reduced.
- During the cultivation of plants such as pistachios, ryaska, eucalyptus and azolla in wastewater, food and space (conditions for laying and reproduction of eggs) are created for fish and other aquatic animals.
- The main areas of animal husbandry are fisheries, poultry, pigs and furs, where biomass of plants such as pistachio, ryaska, eucalyptus and azolla, which are the main food units, can be used.
- The treated wastewater will provide an opportunity to irrigate agricultural crops (cotton, wheat, rice) and get high yields, as well as save water used in agriculture.

As a result of many years of research conducted by world and Uzbek scientists, agricultural enterprises (cattle fattening complexes, poultry) and industrial enterprises (hemp processing, mineral fertilizer production, biochemistry, oil and gas enterprises, silkworm enterprises, textile industry) and utilities - High aquatic plants, pistachios, which use household wastewater from organic-mineral substances, petroleum products and pathogenic microorganisms. ryaska, a new effective biotechnology of biological treatment using eucalyptus and azole has been developed (Shoyoqubov R.Sh.1993.).

In the conditions of Khorezm region, scientific research is being conducted on the use of high aquatic plants in wastewater treatment. Khorezm region has a sharply continental complex climate, the level of salinity is very high. This poses challenges in various areas of agriculture and animal husbandry. The main part of agricultural crops is irrigated. Therefore, the annual water consumption of the region is very high. At the same time, wastewater from communal and agricultural enterprises is added to fresh water through concrete ditches and pipes. This has a negative impact on the ecological environment of the region. Wastewater treatment should be carried out in simple, inexpensive, convenient and environmentally friendly biological methods.

It belongs to the group of high aquatic plants in the regional conditions. Scientists of the Khorezm Mamon Academy are carrying out scientific and

Various animal manure (sheep, cattle, pigs, horses) manure, poultry manure, livestock complexes (poultry factories, pig farms, cattle fattening farms, hemp processing plants, mineral fertilizer plants, biochemical plants) as a nutrient medium for growing pistachios. . ladder factories, meat combines, municipal household (service enterprises)) and sewage canals. Based on the experimental results, an organo-mineral nutrient medium consisting of pig manure, ammonium sulfate, magnesium sulfate and ferric chloride salts is recommended for pistachio breeding. (Kholmurodov et al., 1993).

Results and Discussion

Scientific and practical work has been carried out to increase the number of pistachios in the wastewater of communal and agricultural enterprises of Khorezm region. At the same time, special biological pools have been set up near wastewater treatment plants, agricultural and household waste disposal sites. Sewage was collected in these ponds where pistachios and other biofilter plants were grown, biologically treated, and then discharged or reused. Every ditch, drinking water and sewage was used for planting pistachios. Growing up 25-30% of the crop was harvested daily or every 2-3 days depending on the conditions. A wire mesh strainer was used for collection. In the experimental area of Khorezm Mamun Academy, 3 pools with a depth of 80 cm, a width of 130 cm and a length of 180 cm were created, and 1 pool with a depth of 200 cm and a width of 130 cm and 180 cm was created. Gravel was laid at the bottom, and a series of baked bricks were laid on the sides and insulated. Initially, the first of the 3 biological pools was filled with tap water at a depth of 50 cm, the second with running water, and the third with drainage water, and pistachio bushes were planted and analyzed vegetatively. The artificial biological ponds where pistachios were planted were given a certain amount of organic fertilizer. The pistachio grew well in such conditions. This process indicates that the pistachio is adapted to the conditions of the Khorezm region.

In the next experiment, sewage from Khiva carpets was taken from Khiva carpets and poured into biological pools. The initial physicochemical composition of the wastewater was studied and mixed with drinking water in a ratio of 1: 1, 1: 3, 1: 4. Experiments III option (Option I, Biological Pool 1, 100% Sewage and Pistachio, Option II, Biological Pool 2, sewage + drinking water 1: 1 + pistachio, III-variant, 3-biological pool obucket water + drinking water 1: 3 + pistachio) and 28-day results were recorded. In the experiment, the biomass of the pistachio plant was in the first three days 100% sewage and pistachio in the grown I variant, the wet mass was 1048 g (101%), while at the end of the experiment the total biomass was 2014 g and the daily growth biomass was 168.7 g / m² (201%). Sewage + tap water 1: 1 + pistachio in the grown variant II, the wet mass in the

color changed from yellow to colorless. color, loss of odor, decrease in the amount of suspended solids (107.5 mg / l -72.5 mg / l), increase in the amount of dissolved oxygen in water (1.4 mg / l - 12.3 mg / l), biochemical processes of oxygen for 5-day consumption (KBS5) decreased the amount (123.5 mg / l -28.7 mg / l), the amount of O₂ in the oxidation of organic matter increased from 118.4 mg / l to 28.9 mg / l, and the amount of ammonia 9 , Decreased by 0-1.8 mg / l, as well as the loss of nitrites, nitrates and sulfates 121.0-67.0 mg / l, chlorides 113.0-54.8 mg / l, phosphates 8.3- A decrease of about 1.4 mg / l was noted. (Table 2).

Table 2

Physicochemical composition of wastewater in the pistachio pond (before and after the experiment)

Indicators	Pre-experience situation	Post-experimental condition
Temperature, t°	27.5	33.8
pH	6.8	7.6
Rangi	Reddish-yellow	Colorless
Smell (in points)	5.0	No.
Suspended substances, mg / l	3.8	2.2
Oxygen, mg / l	1.4	12.3
KBS5mg O ₂ / l	123.5	28.7
Oxidation, mg O ₂ / l	118.4	28.9
Ammonia, mg / l	9.0	1.8
Nitrites, mg / l	0.07	No.
Nitrates, mg / l	2.7	No.
Sulfates, mg / l	121.0	67.0
Chlorides, mg / l	113.0	54.8
Phosphates mg / l	8.3	1.4

Conclusion

Pistia during scientific research(Pistia stratiotes).When studying the composition of wastewater in the pond where the plant was grown, the water environment increased from acidic to alkaline, color changed from reddish-yellow to colorless, odorless, removed from biogenic salts, ammonia, nitrite, sulfates, chlorides, photosynthesis, oxygen content in the water the amount of

- Pistia stratiotes* L. for mitigating water pollution in the Egyptian wetlands. *Int J Phyto* 20 (5): 440-449
11. Qutliyev J. (1989). Service of biological pools in wastewater treatment. Tashkent. Uzbekistan. Science Publishing House. 3-12 p.
 12. Laytinen GA, Harris BE. (1979). Chemical analysis. Moscow .s. 395.
 13. Lur'e Yu.Yu. (1984). Analytic chemistry promiyshlenniyx stochnix vod. Moscow, p.446.
 14. Merrill ED (1925). An Enumeration of Philippine Flowering Plants, vol.1 [reprint], 463 pp. Bureau of Printing, Manila (PH).
 15. Muzaffarov AM, Shoyoqubov R.Sh, Yunusov II, Qutliyev J, Abdullayev AA, Khaydarova XN (1983). Opium cultivation *Pistia stratiotes* and using them in cleaning wastewater. *Uz. biologist journal.* №4.s.29-32.
 16. Xayitov Yo.K. (2001). Efficiency of pistachio (*PISTIA STRATIOTES* L) in bioecological treatment of wastewater of spinning and weaving enterprises. Tashkent. Nomzodlik diss. 20-25 b.
 17. Hill MP (2003). The impact and control of alien aquatic vegetation in South African aquatic ecosystems. *African Journal of Aquatic Science*, 28, 19–24.
 18. Hill MP (2013). Report on trip to Morocco for the biological control of *P. stratiotes* at Fez.
 19. Holm LG, Plucknett DL, Pancho JV, Herberger JP. (1977). The world's worst weeds: distribution and biology. Honolulu (HI): The University Press of Hawaii.
 20. Kholmurodov AF, Shoyoqubov R.Sh., Qutliyev J. (1993). Recommendations on the biotechnology of effective wastewater treatment with pistachios. Tashkent. 3-15 p.
 21. Shoyoqubov R.Sh. (1993). Biology of *pistia* and possibilities of their practical usage. Avtoref.doctors.diss. Tashkent. p.46.
 22. Strogonov NS, Buzinova NS (1980). Practical directing on hydrochemistry. Moskva s.194.
 23. Tackholm. V. (1974). Students' Flora of Egypt. 2nd eds. Cairo, Egypt: Cairo Uni. pp. 888.
 24. Waterhouse DF (1997). The major invertebrate pests and weeds of agriculture and plantation forestry in the Southern and Western Pacific, 93 pp. The Australian Center for International Agricultural Research, Canberra, ACT (AU).