

Impact of soil fertilization levels on the activity of invertase and phosphatase enzymes in the soils of Bukhara region

Otabek Umarov^{1,2*}, Zahro Bafayeva^{1,2}, and Dilorom Bo`riyeva¹

¹Bukhara State University, Bukhara, Uzbekistan Street Mukhammad Ikbol, 11, 200118, Uzbekistan

²Bukhara State Pedagogical Institute, Bukhara City, Piridastgir, 2, 200100, Uzbekistan

Abstract. This article provides information on the types and levels of soil fertilization affecting the activity of invertase and phosphatase enzymes in the soils of the Bukhara region. According to the results, the activity of enzymes increases with soil fertilization. It has been scientifically proven that enzymatic activity increases in soils that have not been fertilized, as well as in soils fertilized at moderate and high levels.

1 Introduction

The formation of soil enzymatic activity is closely related to the ecological and biological factors of soil formation. It was found that soils with low organic matter content, poor supply of nutrients, and various degrees of fertilization exhibit weak enzymatic activity [1].

The enzymatic activity in soils has been observed to be relatively higher in all types and layers of soil and tends to decrease towards the lower layers. Studies on the seasonal dynamics of enzymatic activity in soils of the Jizzakh region have shown the importance of hydrothermal conditions and active biochemical processes. Enzymatic activity has been recorded in the upper and gypsum-free horizons of soils with high humus content and in soils with high gypsum content. During the spring season, enzymatic activity was found to be higher in soil layers compared to the summer season. According to the results of microbiological analyses, the gypsum content and degree of soil formation affect the microbiological activity of soils [2,3,4,5,6].

2 Materials and Methods

Scientific research was conducted using expeditionary methods, and soil samples from various types of alluvial soils were collected and analyzed based on genetic horizons. Soil samples based on genetic horizons were taken and enzymatic indicators were analyzed in laboratories. The activity of invertase and phosphatase enzymes was determined using the methods developed by A.Sh.Galstyan [7-15].

* Corresponding author: varzonze1985@mail.ru

3 Results and Discussion

Enzymes participate in the usual oxidation-reduction reactions in the process of oxidation-reduction reactions by soil microorganisms and decomposition of organic matter under normal conditions. This contributes positively to soil fertility, characteristics, and regimes. Humification, hydrolytic processes, synthesis, and other enzymatic effects catalyze in the presence of enzymatic action. Soil enzymes participate in the biochemical processes of organic matter transformation in soil. The identification of enzymes is based on substrate reaction in optimal conditions of temperature, pH (environmental reaction), and substrate concentration at a certain time interval, resulting in the production of a product. For example, polyphenol oxidase catalyzes the conversion of aromatic series organic compounds into soil humus components, while catalyzing the oxidation of phenols to quinones in the presence of atmospheric oxygen.

Hydrolases include invertase, urease, phosphatase, and protease enzymes. The oxidoreductase group includes catalase, peroxidase, polyphenol oxidase, and other enzymes. Peroxidase and polyphenol oxidase enzymes participate in the formation of humus under the influence of atmospheric oxygen and play a crucial role in this process.

The enzyme invertase catalyzes the hydrolytic cleavage of sucrose into glucose and fructose. It also acts on other carbohydrates, producing fructose molecules. According to most researchers, the activity of the invertase enzyme correlates positively with soil fertility level and biological activity compared to other enzymes. The activity of the invertase enzyme indicates the amount of easily hydrolysable carbohydrates in the soil. Easily hydrolysable carbohydrates serve as an energy source for all soil heterotrophs. The activity of the invertase enzyme is influenced by soil fertilization. In unfertilized soil and compared to alluvial soils, the activity of the invertase enzyme was higher. With an increase in fertilization level, the activity of the invertase enzyme decreased. This was particularly noticeable in moderately and highly fertilized alluvial soils. Thus, the sharp increase in salt content in the soil due to irrigation and its high concentration contributed to the positive influence on the activity of the invertase enzyme. With an increase in soil fertilization levels, the harmful and saline salts content, the concentration of sodium, magnesium, and chloride ions, and their proportion increased, which negatively affected the activity of the invertase enzyme. These enzymes also participate in the hydrolytic cleavage of high molecular organic compounds, contributing to the movement of nutrients and organic matter by facilitating their movement through the soil, thereby playing an important role in soil fertility. Soil enzymatic activity is considered an elementary soil classification. This situation is most common in heavily fertilized soils. The activity of the invertase enzyme increased seasonally. This situation is also related to the very active state of the rhizosphere during this period. In addition, the activity of the soil invertase enzyme varied depending on the soil profile. In this case, the activity of the invertase enzyme decreased from the upper layer to the lower layers of the soil. The highest activity was recorded in the upper horizon, while the lowest activity was observed in the lowest layer. The activity of the invertase enzyme was highest in the horizon with the highest amount of humus. The movement of oxygen from the upper horizons to the lower layers of the soil also contributed to the decrease in soil invertase enzyme activity. For example, in unfertilised alluvial soils, during the spring season, the activity of invertase enzyme in the soil layers of 0-25; 25-50; and 50-80 cm was on average 12.5; 10.3; 7.1 mg glucose per g of soil per 24 hours, respectively, while during the summer season, it was found to be 12.2; 10.0; 6.7 mg glucose per g of soil per day, respectively. The soil fertilization level affected the decline in invertase enzyme activity in alluvial soils. For example, in unfertilised alluvial soils, during the spring season, the activity of invertase enzyme in the soil layers of 0-25; 25-50; and 50-80 cm was on average 10.5; 8.3; 6.4 mg glucose per g of soil per day, respectively, while during the

summer season, it was found to be 10.2; 8.0; 6.1 mg glucose per g of soil per day, respectively. The activity of the invertase enzyme was highest in heavily fertilized alluvial soils. For example, in heavily fertilized alluvial soils, during the spring season, the activity of invertase enzyme in the soil layers of 0-25; 25-50; and 50-80 cm was on average 6.3; 4.7; 2.6 mg glucose per g of soil per day, respectively, while during the summer season, it was found to be 6.8; 5.1; 2.9 mg glucose per g of soil per day, respectively. Thus, the soil invertase enzyme activity was influenced by the soil fertilization level, seasonal changes, and genetic horizons in the soil profile.

Table 1. Enzymatic Activity of Alluvial Soil Types at Various Depths of Soil Layer

Soil Horizons, cm	Invertase, 24 hours, mg glucose produced per 1g soil			Phosphatase, 24 hours, mg P ₂ O ₅ produces per 1g soil		
	Seasons of the year					
	Spring	summer	Autumn	Spring	Summer	Autumn
Unsalted						
0-25	12,5	13,8	12,2	9,5	10,1	9,3
25-50	10,3	11,6	10,0	7,2	7,8	7,0
50-80	7,1	7,7	6,7	5,8	6,0	5,5
Weakly salted						
0-25	10,5	11,1	10,2	9,0	9,7	8,8
25-50	8,3	9,6	8,0	7,9	8,4	7,5
50-80	6,4	7,0	6,1	6,1	6,6	5,7
Moderately salted						
0-25	7,2	7,6	7,0	6,7	7,0	6,5
25-50	5,3	5,8	5,0	5,0	5,8	4,8
50-80	3,5	3,9	3,1	3,1	3,5	2,9
Strongly salted						
0-25	6,3	6,8	6,1	5,9	6,3	5,8
25-50	4,7	5,1	4,4	4,4	5,0	4,2
50-80	2,6	2,9	2,3	2,2	2,6	2,0
S _x %	2,99	1,93	2,62	4,79	3,97	4,22
NSR _{0,5}	0,61	0,43	0,51	0,84	0,75	0,64

Phosphatase is one of the important enzymes. Soil can contain a significant amount of phosphorus in organic forms. Microorganisms possessing the specific enzyme phosphatase catalyze the hydrolysis of orthophosphate esters, releasing phosphoric acid. The general activity of phosphatase in the soil depends on the amount of humus and organic phosphorus. Organic phosphorus serves as a substrate for the phosphatase enzyme. Phosphatase (both alkaline and acid) catalyzes the hydrolysis of a variety of phosphoric organic compounds into orthophosphates. The activity of phosphatase in the soil is higher when the phosphorus content in the soil is lower. Therefore, the activity of soil phosphatase can be used as an additional indicator to determine the need for the use of phosphorus-containing fertilizers in the soil. The highest activity of soil phosphatase is observed in the

rhizosphere of plants. The activity of soil phosphatase enzyme changes depending on the degree of soil cultivation.

The activity of phosphatase enzyme in unsalted alluvial soil has increased compared to moderately salted alluvial soil. With the increase in cultivation intensity, the activity of soil phosphatase has gradually increased. In this case, alluvial soil with moderate and strong cultivation intensity showed strong manifestations. For example, in unsalted alluvial soil, the activity of phosphatase enzyme in the spring at depths of 0-25; 25-50; and 50-80 cm was found to be 9.5; 7.2; 5.8 mg P₂O₅ /g soil per 24 hours, respectively, while in the summer, it was found to be 10.1; 7.8; 6.0 mg P₂O₅ /g soil per 24 hours, respectively. The beginning of soil cultivation coincides with the onset of phosphatase enzyme activity.

In addition to soil cultivation, seasonal variations, the level of crop development, and the depth of soil layers significantly affect the activity of phosphatase enzymes. In the spring, the activity of phosphatase enzyme in the soil was highest. In the summer, the activity of the phosphatase enzyme slightly increased compared to the spring. These changes are also associated with soil moisture. When moisture levels increase sharply, the activity of the phosphatase enzyme increases. This increase is also caused by the concentration of salts, harmful and soluble salts, magnesium, sodium, and chloride ions. They negatively affect soil phosphatase activity. For example, in moderately cultivated alluvial soil, the activity of phosphatase enzyme in the spring at depths of 0-25; 25-50; and 50-80 cm was found to be 6.7; 5.0; 3.1 mg P₂O₅ /g soil per 24 hours, respectively, while in the summer, it was found to be 7.0; 5.8; 3.5 mg P₂O₅ /g soil per 24 hours, respectively. In strongly cultivated alluvial soil, the activity of the phosphatase enzyme was the lowest. This is associated with the concentration of salts, harmful and soluble salts, magnesium, sodium, and chloride ions, as well as their maximum accumulation. For example, in strongly cultivated alluvial soil, the activity of phosphatase enzyme in the spring at depths of 0-25; 25-50; and 50-80 cm was found to be 5.9; 4.4; 2.2 mg P₂O₅ /g soil per day, respectively, while in the summer, it was found to be 6.3; 5.0; 2.6 mg P₂O₅ /g soil per day, respectively. Therefore, soil phosphatase activity is closely related to soil cultivation, the content of harmful and soluble salts, magnesium, sodium, and chloride ions, soil layer depth, seasonal changes, and the development period of crops.

4 Conclusion

The cultivation of alluvial soils has a positive effect on their enzymatic activity. In this regard, the activity of hydrolytic enzymes such as invertase and phosphatase significantly increases. These cultivated soils contribute to the rapid decomposition of organic matter, ammonification, and oxidation-reduction processes.

References

1. Kh. Artikova, S. Shadiyeva. Ecological condition of soils, problems of pollution and its prevention. *E3S Web of Conferences* **389**, 04014 (2023)
2. Hamroyev, H. Jumayeva, *E3S Web of Conferences* **420**. 10007 (2023)
3. A.Ilyosoy. The influence of sowing rates of winter wheat varieties planted in Bukhara region on seed viability and actual seedling thickness. *E3S Web of Conferences* **462**, 02001 (2023)
4. D. Bo'riyeva, M. Xalilova, and Z.Atayeva. Dependence of microbiological activity of alluvial soils of irrigated meadow of Bukhara oasis on salinity levels. *E3S Web of Conferences* **420**, 03013 (2023)

5. B. Mamurov et al., E3S Web of Conferences **538**. 05025 (2024)
6. B. Mamurov et al., E3S Web of Conferences **538**. 05031 (2024)
7. H. Salimova and H. Artikova. Determination of the mechanical composition and salinity of the Bukhara Gijduvan district of the Bukhara region E3S Web of Conferences **389**, 04012 (2023)
8. M. Sattorova, and M.Raximova The mechanical composition of irrigated sandy desert soils in the Jondor district of Bukhara region, and the amount of nutrients in them. E3S Web of Conferences **420**, 03010 (2023)
9. M. Shirinova et al., E3S Web of Conferences **538**. 05016 (2024)
10. R. Kuldoshev et al., E3S Web of Conferences **371**. 05069 (2023)
11. R. Qo'ldoshev et al., E3S Web of Conferences **538**. 05017 (2024)
12. R. Qo'ldoshev et al., E3S Web of Conferences **538**. 05042 (2024)
13. S.M.Nazarova. Factors of soil formation and their evolution in the south of the Zerafshan valley. E3S Web of Conferences **389**, 04016 (2023)
14. U. Khayitov et al., E3S Web of Conferences **538**. 05022 (2024)
15. G. Akramova et al., E3S Web of Conferences **538**. 05034 (2024)