

OPEN ACCESS  
PEER REVIEWED

Volume 2, Issue 4  
April, 2024

ISSN (E): 2938-379X  
IFSIJ JIF: 8.325  
SJIF 2024: 5.178

# WEB OF TEACHERS

INDERSCIENCE RESEARCH



 <https://webofjournals.com>

 [editor@webofjournals.com](mailto:editor@webofjournals.com)



WEB OF JOURNALS  
OPEN ACCESS JOURNALS PUBLISHING

## THE ROLE OF IMMUNOSTIMULANTS IN INCREASING THE RESISTANCE OF COTTON TO GUMMOSIS

R. O. Atoeva  
G. A. Goziyeva,  
S. S Bakayeva

Bukhara State University  
r.o.atoyeva@buxdu.uz

### Abstract

In this article, information about the role of the immunostimulants Fitovak and Zerox on the gummosis disease of the cotton plant. The obtained results showed that Zerox immunostimulant played an important role in increasing the resistance of cotton to gummosis compared to Fitovak immunostimulant.

**Keywords:** Bukhara-10, Zerox, Fitovak, gummosis, disease, biological efficiency.

### Introduction

Immunostimulants and stimulants protect plants from biotic and abiotic influences, reduce stress-causing reactive oxygen species, activate the antioxidant defense system in them, or create an immune system in plants by increasing salicylic acid compounds [1]; [2], [3].

To give plants resistance to diseases, one of the important factors is the use of various immunostimulants in order to form resistant forms [4]. Here is one such immunostimulant - the drug Zerox of universal and contact action, containing 3000 mg/l of silver ions. Silver ions at a concentration of 3000 mg/l contained in this preparation are the first to penetrate the membranes of fungi and bacteria parasitizing the plant and destroy them. Secondly, by interacting with plant receptors and enzymes, it produces high levels of oxygen in the plant cell and creates systemic immunity against various pathogens in the plant body.

With gummosis, fluid leaks out as a result of melting of the skin parenchyma in the internal tissue of the tuber. To recognize this disease, it is necessary to know the relationship between the fruit of the "host plant" (cotton) and the "parasite" [5].

Studies have been conducted on the effect of this drug on gummosis disease of the cotton variety Bukhara-10 under field conditions.

### Research Methodology

To determine the effectiveness of the drugs applied to the seeds during the research, diseases of root rot and gummosis were determined in sprouted seedlings according to the Manners scale and calculated using the Abbott formula [6].

In order to clarify the spectrum of action of the drug, research work was carried out in the conditions of moderately saline soils of the pilot farm of PSUAITI "Bukhara". As a result, it was established what dose (1.0-2.0-3.0 l/ha) of the Zerox immunostimulant provides high resistance to gummosis. According to the experimental system, all options for using the Zerox immunostimulant were compared with the standard immunostimulant (Fitovak 200 ml/t, ml/ha) and the control option (drinking water).

### Analysis and Results

In the years under study, before planting seeds in the same sowing season, there was a lot of precipitation, which somewhat delayed the timing of planting seeds, but due to the low amount of precipitation after sowing and dry and hot air temperatures, the number of infections of cotton seedlings with gummosis in all variants was significantly less, and in experimental variants it was practically not observed. Even in the control variants, the symptoms of this disease were very mild (Table 1).

Table 1 Influence of immunostimulants in increasing the resistance of cotton to gummosis

№	Name of the drug	The number of seedlings	Incidence of gummosis disease by years during the growing seasons of cotton				
			on gummosis disease				
			2018	2019	2020	an average of three years	BC,%
The number of infected sprouts in 90m <sup>2</sup> after the drug was applied during the ripening period, pcs							
1	Control	80-90	5	6	3	4,7	0
2	Control	100-110	6	7	5	6,0	0
3	Control	120-130	6	8	7	7,0	0
4	Fitovak-200 ml/t; l/ha	80-90	0	0	0	0,0	100,0
5	Fitovak-200 ml/t; l/ha	100-110	0	1	0	0,3	93,6
6	Fitovak-200 ml/t; l/ha	120-130	0	2	1	1,0	85,7
7	Zerox 1 l/t, l/ha	80-90	0	0	0	0,0	100,0
8	Zerox 1 l/t, l/ha	100-110	0	1	0	0,3	93,6
9	Zerox 1 l/t, l/ha	120-130	0	2	1	1,0	85,7
10	Zerox 2 l/t, l/ha	80-90	0	0	0	0,0	100,0
11	Zerox 2 l/t, l/ha	100-110	0	0	0	0,0	100,0
12	Zerox 2 l/t, l/ha	120-130	0	0	0	0,0	100,0
13	Zerox 3 l/t, l/ha	80-90	0	0	0	0,0	100,0
14	Zerox 3 l/t, l/ha	100-110	0	0	0	0,0	100,0
15	Zerox 3 l/t, l/ha	120-130	0	0	0	0,0	100,0

According to the scientific results obtained, it was established that the number of infected seedlings was less in areas (90 m<sup>2</sup>) of the tested options during the period of 2-4 true leaves, in options with in the number seedling of 80-90 thousand plant/ha compared to other seedling thicknesses.

Control, standard and Zerox 1-2-3 l/t, where, when the cotton ripened, in the number seedling of 80-90 thousand hectares; 1-2-3 l/ha + surfactant 0.15 l/ha were used in options 1-4-7-10-13, after using the drug the average number of infected seedlings over three years: 4.7-0.0-0, 0-0.0-0.0 units, the number of seedlings infected with gummosis in these variants with a standing density of 100-110 thousand hectares, respectively: 6-0.3-0.3-0.0-0.0 units, in those variants with a in the number seedlings of 100-110 thousand hectares, the number of infected seedlings was: 7-1.0-1.0-0.0-0.0 pcs.

Thus, according to the scientific results obtained over 3 years, during the initial and ripening periods of cotton, with a standing density of 80-90 thousand hectares, the Zerox preparation is applied at the rate of 2 l/t per 1 hectare. for seedlings and 2 l/ha + surfactant 0.15 l/during the cotton growing season. In the 10th option, the highest BS by phase was 100 - 100%.

From the period of 2-4 leaves of cotton to the ripening period, the parameters for the prevention of gummosis disease on an area of 90 m<sup>2</sup> are 80-90 thousand plant/ha of in the number seedling and 1-3 l/t of Zerox. Options using rates of 1-3 l/ha + surfactant 0.15 l/ha showed intermediate results.

Depending on the increase in the number of the seedlings in the standard and Zerox 1l/ha norms, a decrease in the optimal the number seedlings was observed by 14.3-14.3% in the BS ripening phase compared to the options where the number of the seedlings was left at the norm and norm duration was used.

### Conclusion

Another of the most effective ways to increase the resistance of cotton to gummosis is to depilate the seeds and treat them with Zerox at a rate of 2 l/t before planting and 2 l/ha + surfactant 0.15 l. /ha during the period of plant growth - foliar feeding.

### References

1. Икромова М.Л. Тупроқ унудорлигини сақлаш ва гўза вилтига қарши курашда «Фитовак» ва микробиологик препаратларни композицион ҳолда қўллашнинг аҳамияти // Тупроқшунослик-мамлакат экологик ва озиқ- овқат хавфсизлиги хизматида” Республика илмий-амалий анжумани. Тошкент, 2017.-Б. 59-63.
2. Atoeva, R. O., & Hotamova, H. (2023). Improvement of valuable signs by using “Zerox” immunostimulant in cotton. In E3S Web of Conferences (Vol. 389, p. 03114). EDP Sciences.
3. Ikramova, M. L., and R.O. Atoeva. "INFLUENCE OF ZEROX IMMUNOSTIMULANT ON THE GROWTH AND DEVELOPMENT OF COTTON." ILMIIY XABARNOMA 196: 36.
4. Ikramova, Mahbuba Latibovna, Rukhsora Odilovna Atoeva, and Dilsora Odilovna Atoeva. "Influence of the Zerox Immunostimulant on Cotton Production." American Journal of Plant Sciences 11.4 (2020): 564-568.

5. Маннанов Р.Н. Взаимоотношения почвенных антогонистов с некоторыми фитопатогенами, вызывающими основные болезни хлопчатника (гоммоз, корневая гниль, фузориоз) и пшеницы (корневая гниль). Автореф. – Тошкент – 2010. – б. 5-24.
6. Manners J.G. Studies on the physiologic specialization of yellow rust (*Puccinia glum arum* [Schmidt] Erik's) in Great Britain. *Annals of Applied Biology*.1950, 37, 2: p -187-214.



25. SPECIFIC CHARACTERISTICS OF DOUBLE FERTILITY SIMMENTAL CATTLE (COWS) IMPORTED TO KARAKALPAKSTAN  
Naurizov Tokhtamis Kidirbergenovich, Kazakbaev Bakhit Seylbekovich, Kutlimuratov Maman  
125-127
26. THE ROLE OF IMMUNOSTIMULANTS IN INCREASING THE RESISTANCE OF COTTON TO GUMMOSIS  
R. O. Atoeva, G. A. Goziyeva, S. S Bakayeva  
128-131
27. PREPARATION OF FLOORS IN EARTHQUAKE AREAS AND DESIGN OF LOAD-BEARING STRUCTURES IN EARTHQUAKE-PROBENT AREAS  
Teshaboeva N. D., Kasimova H., Tursunova S.M.  
132-136
28. POLITICAL RULE OF THE CUSHAN DYNASTY CAPTURE  
Nuriddin Fatxullayev  
137-139
29. DEVELOPING STUDENTS' PHYSICAL QUALITIES IN THE TRAINING PROCESS WHILE USING GAMES IN BOXING SPORTS  
Rixsiyev Dilshod Shavkatovich  
140-143
30. CHARACTERISTICS OF LEXICAL STYLISTIC METHODS IN THE WORK OF ANDISHA AND PRIDE  
Panjiyeva Nasiba, Ibragimova Lobar Doniyorovna  
144-146
31. BUSINESS MANAGEMENT AND ITS INDIVIDUAL  
Ergashev Zafar Alibekovich  
147-150
32. THE INFLUENCE OF THE WORLD'S FOOD TRADITIONS ON ATHEROSCLEROSIS REVIEW  
Дон А. Н., Гиясова Н. А., Шагулямова К. Л., Мирзахмедова Н. А.  
151-158
33. IMPROVING THE PROCEDURE FOR ORGANIZING STATE FINANCIAL CONTROL  
Khonnazarov Islom Shodievich  
159-161
34. NAVIGATING DIVERSITY: A COMPREHENSIVE EXPLORATION OF INCLUSIVE TEACHING STRATEGIES IN FOREIGN LANGUAGE EDUCATION IN UZBEKISTAN  
Fayzieva Nilufar  
162-169
35. FAMILIARIZATION WITH INFORMATION ABOUT MODERN BIOINFORMATIC DATABASES  
Amonov Abdullaziz Uktamovich  
170-173
36. METHODS OF DISTINGUISHING PHYTONEMATODES IN PLANTS  
E. Q. Abduraxmonova  
174-176
37. FEATURES OF TRANSLATING METAPHORICAL PHRASES IN POLITICAL TEXTS INTO UZBEK AND ENGLISH  
Abdumuratova Farangiz Alisherovna  
177-184